

Examples of the Iwasawa Invariants and the Higher K -groups Associated to Quadratic Fields

Dedicated to Professor Toru Ishihara on his 65th birthday

By
Hiroki SUMIDA-TAKAHASHI

*Faculty and School of Engineering
The University of Tokushima
Minami-josanjima, Tokushima 770-8506, JAPAN
e-mail address: hiroki@pm.tokushima-u.ac.jp*
(Received September 28, 2007)

Abstract

We compute the Iwasawa invariants of $\mathbf{Q}(\sqrt{f}, \zeta_p)$ in the range $|f| < 200$ and $5 \leq p < 200000$ (resp. $|f| < 10$ and $5 \leq p < 1000000$). These computational results give us concrete information on the higher K -groups of the ring of integers of $\mathbf{Q}(\sqrt{f})$.

2000 Mathematics Subject Classification. 11R23, 11R70

Introduction

Let F be a number field and \mathcal{O}_F the ring of integers of F . Put $K = F(\zeta_p)$ and denote by K_∞ the cyclotomic \mathbf{Z}_p -extension of K . Let L_∞ be the maximal unramified abelian p -extension of K_∞ and L'_∞ the maximal unramified abelian p -extension of K_∞ in which every prime divisor lying above p splits completely. Put $X_\infty = \text{Gal}(L_\infty/K_\infty)$ and $X'_\infty = \text{Gal}(L'_\infty/K_\infty)$.

It is known that there are relations between Iwasawa modules X'_∞ and Quillen's K -groups $K_n(\mathcal{O}_F)$. The main purpose of this paper is to give concrete information on the Iwasawa invariants of X_∞ and the higher K -groups $K_n(\mathcal{O}_F)$ for quadratic fields F by using these relations.

Following [9, 10], we compute Iwasawa invariants and found some exceptional pairs. Using these pairs, we give exceptional examples of $K_n(\mathcal{O}_F)$. For example, we find that for $5 \leq p < 1000000$, p divides the order of $K_{33588}(\mathcal{O}_{\mathbf{Q}(\sqrt{8})})$ if and only if $p = 7$ or 157229 under the Quillen-Lichtenbaum conjecture.

1 Iwasawa invariants of $\mathbf{Q}(\sqrt{f_\chi}, \zeta_p)$

Let χ be a quadratic Dirichlet character and p an odd prime number. Assume that $\chi \neq \omega^{\frac{p-1}{2}}$, where $\omega = \omega_p$ is the Teichmüller character $\mathbf{Z}/p\mathbf{Z} \rightarrow \mathbf{Z}_p$ such that $\omega(a) \equiv a \pmod{p}$. Put $F = F_\chi = \mathbf{Q}(\sqrt{f_\chi})$ and $K_n = \mathbf{Q}(\sqrt{f_\chi}, \zeta_{p^{n+1}})$. Let A_n (resp. A'_n) the p -part of the ideal class group (resp. p -ideal class group) of K_n .

Put $G_\infty = \text{Gal}(K_\infty/F)$, $\Delta = \text{Gal}(K_\infty/F_\infty)$ and $\Gamma = \text{Gal}(K_\infty/K)$. Further put $\Delta' = \text{Gal}(K_\infty/\mathbf{Q}_\infty)$ and $e'_\psi = \frac{1}{\#\Delta'} \sum_{\delta \in \Delta'} \psi(\delta) \delta^{-1}$ for a Dirichlet character ψ of Δ' . For a $\mathbf{Z}_p[\Delta']$ -module A , we denote $e'_\psi A$ by A^ψ . Let $\lambda_p(\psi)$, $\mu_p(\psi)$ and $\nu_p(\psi)$ (resp. $\lambda'_p(\psi)$, $\mu'_p(\psi)$ and $\nu'_p(\psi)$) be the Iwasawa invariants associated to X_∞^ψ (resp. X'_∞^ψ), i.e.,

$$\#A_n^\psi = p^{\lambda_p(\psi)n + \mu_p(\psi)p^n + \nu_p(\psi)} \quad (\text{resp. } \#A'_n^\psi = p^{\lambda'_p(\psi)n + \mu'_p(\psi)p^n + \nu'_p(\psi)})$$

for sufficiently large n . By Ferrero-Washington's theorem, we have $\mu_p(\psi) = \mu'_p(\psi) = 0$ for all p and ψ .

Assume that ψ is even. The Iwasawa polynomial $g_\psi(T) \in \mathbf{Z}_p[T]$ for the p -adic L -function is defined as follows. Let $L_p(s, \psi)$ be the p -adic L -function constructed by [6]. Let f_0 be the least common multiple of f_ψ and p . By [3, §6], there uniquely exists $G_\psi(T) \in \mathbf{Z}_p[[T]]$ satisfying

$$G_\psi((1 + f_0)^{1-s} - 1) = L_p(s, \psi)$$

for all $s \in \mathbf{Z}_p$ if $\psi \neq \chi^0$. By [2], it was proved that p does not divide $G_\psi(T)$. Therefore, by the p -adic Weierstrass preparation theorem, we can uniquely write

$$G_\psi(T) = g_\psi(T)u_\psi(T),$$

where $g_\psi(T)$ is a distinguished polynomial of $\mathbf{Z}_p[T]$ and $u_\psi(T)$ is an invertible element of $\mathbf{Z}_p[[T]]$. Put $\tilde{\lambda}_p(\psi) = \deg g_\psi(T)$.

For a pair (p, ψ) , we assume the following condition

$$(C) \quad \psi(p) \neq 1 \text{ and } \psi^{-1}\omega(p) \neq 1.$$

If $\psi(p) \neq 1$, we have $\lambda_p(\psi) = \lambda'_p(\psi)$ and $\nu_p(\psi) = \nu'_p(\psi)$.

We extend the tables of [9, 10] to all primes below 200000.

Proposition 1 *For $|f| < 200$ and $100000 < p < 200000$, all exceptional pairs $(p, \chi\omega^k)$ are given in the following table. The meaning of the symbols are as follows: $[\nu] : \nu(\chi\omega^k) > 0$, $[a_0] : v_p(a_0) > 1$, $[b_0] : v_p(b_0) > 1$, $[\text{lmd}] : \tilde{\lambda}(\chi\omega^k) > 1$, where $a_0 = L_p(1, \chi\omega^k)$ and $b_0 = L_p(0, \chi\omega^k)$.*

Exceptional pairs $(p, \chi\omega^k)$ for $100000 < p < 200000$

f	p	k	f	p	k	f	p	k
	[ν]			[a_0]				[lmd]
8	157229	140434	57	119627	53592	133	185189	119132
28	109829	45474	77	175843	43682	168	104971	21988
56	100937	93200	141	120823	39250	-187	166823	150305
-79	153059	68171	157	150401	101272			
-91	107449	81489		[b_0]				
-104	184157	53783	-19	165667	11685			
-119	112241	37701	53	167593	99386			
-120	126691	28093	-71	177473	58993			
149	109211	11960	137	124493	41762			
-183	104803	58845	-152	104399	90165			

Proposition 2 For $|f| < 10$, i.e., $f = -3, 5, -4, -7, 8$ or -8 and $200000 < p < 1000000$, there is only one exceptional pair $(399181, \chi_{-4}\omega^{1683})$, which satisfies $\tilde{\lambda}(\chi_{-4}\omega^{1683}) > 1$.

In Figures 1-2, we compare the actual number of exceptional pairs with the expected number E in the range $200 < p < 200000$.



FIGURE 1. Exceptional pairs (quadratic, $1 < |f| < 200$, $200 < p < 200000$)

FIGURE 2. Exceptional pairs (Quadratic, $1 < |f| < 200$, $200 < p < 200000$)

From our data, the actual numbers still seem to be near to the expected numbers.

2 Higher K -groups of \mathcal{O}_F

We recall some results on Quillen's K -groups.

Theorem 1 (Quillen) *For all $n \geq 0$, $K_n(\mathcal{O}_F)$ is a finitely generated \mathbf{Z} -module.*

Theorem 2 (Borel) *For $m \geq 1$,*

$$\text{rank}_{\mathbf{Z}}(K_{2m-1}(\mathcal{O}_F)) = \begin{cases} r_1(F) + r_2(F) & \text{if } m \text{ is odd,} \\ r_2(F) & \text{if } m \text{ is even,} \end{cases}$$

where $r_1(F)$ is the number of real embeddings of F , and $r_2(F)$ is the number of pairs of complex embeddings of F . Further,

$$K_{2m-2}(\mathcal{O}_F) \text{ is finite.}$$

Conjecture 1 (The Quillen-Lichtenbaum conjecture) *The natural map (via p -adic Chern characters)*

$$K_{2m-i}(\mathcal{O}_F) \otimes \mathbf{Z}_p \rightarrow H_{\text{ét}}^i(\text{Spec}(\mathcal{O}_F[1/p]), \mathbf{Z}_p(m))$$

is an isomorphism for all $m \geq 2$, $i = 1, 2$ and any odd prime number p , where $A(m)$ is the m -th Tate twist of a Galois module A .

The surjectivity of p -adic Chern characters was proved by [1, 4, 7, 8]. We simply denote $H_{\text{et}}^i(\text{Spec}(\mathcal{O}_F[1/p]), A)$ by $H^i(\mathcal{O}_F, A)$.

Theorem 3 ([5, §3, §4]) *For $m \neq 0$, we have*

$$H^1(\mathcal{O}_F, \mathbf{Z}_p(m))_{\text{tors}} \simeq H^0(\mathcal{O}_F, \mathbf{Q}_p/\mathbf{Z}_p(m)).$$

For $m \neq 1$, we have an exact sequence

$$\begin{aligned} 0 \rightarrow X'_\infty(m-1)_{G_\infty} &\rightarrow H^2(\mathcal{O}_F, \mathbf{Z}_p(m)) \\ &\rightarrow \prod_{v|p} H^2(F_v, \mathbf{Z}_p(m)) \rightarrow H^0(\mathcal{O}_F, \mathbf{Q}_p/\mathbf{Z}_p(1-m))^\vee \rightarrow 0, \end{aligned}$$

where $A^\vee = \text{Hom}_{\mathbf{Z}_p}(A, \mathbf{Q}_p/\mathbf{Z}_p)$.

From now on, we use the same notation as in the previous sections. For an even character $\chi\omega^{1-m}$, we write the Iwasawa polynomial $g_{\chi\omega^{1-m}}(T)$ for the p -adic L -function $L_p(s, \chi\omega^{1-m})$ in the form

$$g_{\chi\omega^{1-m}}(T) = \prod_{i=1}^{\tilde{\lambda}(\chi\omega^{1-m})} (T - \alpha_{\chi\omega^{1-m}, i}), \quad \alpha_{\chi\omega^{1-m}, i} \in \overline{\mathbf{Q}_p}.$$

We put

$$x(p, \chi, m-1) = \min\{\nu_p(\chi\omega^{1-m}), v_p(\prod_{i=1}^{\tilde{\lambda}} (1 - (1 + f_0)^{m-1}(\alpha_{\chi\omega^{1-m}, i} + 1)))\}.$$

For an odd character $\chi\omega^{1-m}$, we put $\alpha_{\chi\omega^m, i}^* = \frac{f_0 - \alpha_{\chi\omega^m, i}}{1 + \alpha_{\chi\omega^m, i}}$,

$$g_{\chi\omega^m}^*(T) = \prod_{i=1}^{\tilde{\lambda}(\chi\omega^m)} (T - \alpha_{\chi\omega^m, i}^*)$$

and

$$x^*(p, \chi, m-1) = v_p(\prod_{i=1}^{\tilde{\lambda}(\chi\omega^m)} (1 - (1 + f_0)^{m-1}(\alpha_{\chi\omega^m, i}^* + 1))).$$

Further, for an integer m , we define the following sets of prime numbers

$$\begin{aligned} S_1(\chi, m-1) &= \{p : p'|(m-1), (p-1) \nmid (m-1), \chi\omega^{p'}(p) = 1, \chi\omega^{p'} \neq \chi^0\}, \\ S_2(\chi, m-1) &= \{p : (p-1)|(m-1), \chi(p) = 1\}, \end{aligned}$$

where $p' = \frac{p-1}{2}$. We put

$$y(p, \chi, m-1) = \begin{cases} v_p(m-1) + 1 & \text{if } p \in S_1(\chi, m-1) \cup S_2(\chi, m-1), \\ 0 & \text{otherwise.} \end{cases}$$

Proposition 3 Let χ be a quadratic Dirichlet character, p an odd prime number and $F = F_\chi$. For an odd character $\chi\omega^{1-m}$, if $(p, \chi\omega^m)$ satisfies (C), then

$$\#X'_\infty(m-1)_{G_\infty}^\chi = p^{x^*(p, \chi, m-1)}.$$

For an even character $\chi\omega^{1-m}$, assume that $X'_\infty^{\chi\omega^{1-m}}$ is finite. If $(p, \chi\omega^{1-m})$ satisfies (C) and if $g_{\chi\omega^{1-m}}(T)$ is an Eisenstein polynomial or of degree one, then

$$\#X'_\infty(m-1)_{G_\infty}^\chi = p^{x(p, \chi, m-1)}.$$

Further, for an integer m , we have

$$\frac{\#\prod_{v|p} H^2(F_v, \mathbf{Z}_p(m))^\chi}{\#H^0(\mathcal{O}_F, \mathbf{Q}_p/\mathbf{Z}_p(1-m))^\chi} = p^{y(p, \chi, m-1)}.$$

Proof. In [9, Proposition 4.1], we prove the above theorem when χ is even. In the same way, using an isomorphism

$$X'_\infty(m-1)_\Delta^\chi \simeq X'_\infty^{\chi\omega^{1-m}} \otimes \mathbf{Z}_p(m-1),$$

we can show the above equations. \square

For a positive integer m and a prime number p , we denote by $K'_{2m-2}(\mathcal{O}_F)(p)$ the p -Sylow subgroup of $K_{2m-2}(\mathcal{O}_F)$. Here we put

$$\begin{aligned} K'_{2m-2}(\mathcal{O}_F) &= \bigoplus_{5 \leq p < 200000} K_{2m-2}(\mathcal{O}_F)(p), \\ X'(\chi, m-1) &= \prod_{5 \leq p < 200000} \#X'_\infty(m-1)_{G_\infty}^\chi \text{ and} \\ Y'_i(\chi, m-1) &= \prod_{p \in S_i(\chi, m), 5 \leq p < 200000} \frac{\#\prod_{v|p} H^2(F_v, \mathbf{Z}_p(m))^\chi}{\#H^0(\mathcal{O}_F, \mathbf{Q}_p/\mathbf{Z}_p(1-m))^\chi}. \end{aligned}$$

Then, Theorem 3 and the surjectivity of p -adic Chern characters, we have

$$\#K'_{2m-2}(\mathcal{O}_F)^\chi \text{ is divided by } X'(\chi, m-1) \cdot Y'_1(\chi, m-1) \cdot Y'_2(\chi, m-1).$$

For an odd character $\chi\omega^{1-m}$, we can compute $v_p(X'(\chi, m-1))$ from the zeros of the Iwasawa polynomial by Proposition 3. In fact, we can easily obtain a lot of examples of (χ, m) with $X'(\chi, m-1) > 1$.

On the other hand, for an even character $\chi\omega^{1-m}$, it is more difficult to obtain examples of (χ, m) with $X'(\chi, m-1) > 1$. Since Vandiver's conjecture is true for all $p < 12000000$, $X'_\infty(m-1)_{G_\infty}^{\chi^0}$ is trivial for any odd integer m . Further we have $\#H^2(\mathbf{Q}_p, \mathbf{Z}_p(m)) = \#H^0(\mathbf{Q}_p, \mathbf{Q}_p/\mathbf{Z}_p(1-m)) = \#H^0(\mathbf{Z}, \mathbf{Q}_p/\mathbf{Z}_p(1-m))$. By Proposition 3 and our computational result [9, 10], we obtain such examples in the following tables.

Factors of $\#K'_{2m-2}(\mathcal{O}_{\mathbf{Q}(\sqrt{f_\chi})})$ with $X' = X'(\chi, m-1) > 1$

$-200 < f_\chi < 0$ and $5 \leq p < 200000$

$2m - 2$	f_χ	X'	$2m - 2$	f_χ	X'
122	-4	379	46	-11	79
22	-11	173	5470	-15	4909
38	-19	37	58	-19	41
594	-19	2251	1714	-20	20261
34	-23	193	30	-31	131
1090	-31	821	26	-40	97
198	-51	557	5918	-51	6553
78178	-55	41189	46	-67	433
26	-71	17	14	-79	17
55534	-79	45943	169774	-79	153059
654	-84	10133	47958	-88	33049
30	-91	37	7550	-91	7069
51918	-91	107449	26	-103	17
102	-103	67	35102	-104	17837
260746	-104	184157	1034	-116	4363
149078	-119	112241	3846	-120	4177
197194	-120	126691	26	-127	67
1450	-131	853	14	-136	11
96258	-136	54547	4434	-139	4451
18	-148	23	490	-152	863
1398	-152	3019	6478	-155	12377
46	-163	79	1102	-167	797
91914	-183	104803	30	-187	79

We have $Y'_1(\chi, m-1) = Y'_2(\chi, m-1) = 1$ for all the above cases.

$1 < f_\chi < 200$ and $5 < p < 200000$

$2m - 2$	f_χ	X'	Y'_2	$2m - 2$	f_χ	X'	Y'_2
68372	8	34301	1	33588	8	157229	7
316	12	701	1	96	21	199	5·17
128708	28	109829	47	44	33	53	1
20	37	43	11	936	53	1033	$7 \cdot 13^2 \cdot 37$
15472	56	100937	5	92652	56	55621	43·6619·15443
8	69	19	5	1220	85	3697	1
88	88	71	1	5124	101	5333	43·367
8	104	19	5	20	113	43	11
3540	113	3373	7·11·31	140	124	197	11
380	124	239	11	76	129	67	1
9260	140	4751	1	1208	141	5431	5
20	149	43	1	108	149	71	7·19
92	149	229	47	194500	149	109211	251
90936	156	50051	5·7·19	688	157	401	173
156	161	101	1	28	168	37	1
124	172	73	1	4	173	7	1
20	173	43	1	116	173	101	1
20	177	17	11	36	181	71	1
10724	181	6991	1	944	185	827	1
2904	188	1621	23·67·727	11380	193	62791	1
296	197	521	1				

We have $Y'_1 = Y'_1(\chi, m - 1) = 1$ for all the above cases.

Examples

There exist submodules A_i of K -groups such that

$$K_{122}(\mathcal{O}_{\mathbf{Q}(\sqrt{-4})}) \supseteq K'_{122}(\mathcal{O}_{\mathbf{Q}(\sqrt{-4})}) \supseteq A_1 \simeq \mathbf{Z}/(379\mathbf{Z}),$$

$$K_{68372}(\mathcal{O}_{\mathbf{Q}(\sqrt{8})}) \supseteq K'_{68372}(\mathcal{O}_{\mathbf{Q}(\sqrt{8})}) \supseteq A_2 \simeq \mathbf{Z}/(34301\mathbf{Z}), \text{ and}$$

$$K_{33588}(\mathcal{O}_{\mathbf{Q}(\sqrt{8})}) \supseteq K'_{33588}(\mathcal{O}_{\mathbf{Q}(\sqrt{8})}) \supseteq A_3 \simeq \mathbf{Z}/(7 \cdot 157229\mathbf{Z}).$$

References

- [1] W. Dwyer and E. Friedlander, *Algebraic and étale K -theory*, Trans. Amer. Math. Soc. **292** (1985), 247–280.
- [2] B. Ferrero and L. Washington, *The Iwasawa invariant μ_p vanishes for abelian number fields*, Ann. of Math. **109** (1979), 377–395.
- [3] K. Iwasawa, *Lectures on p -adic L -functions*, Ann. of Math. Stud., vol. 74, Princeton Univ. Press: Princeton, N.J., 1972.
- [4] B. Kahn, *On the Lichtenbaum-Quillen conjecture*, vol. 407, pp. 147–166, Algebraic K -theory and algebraic topology (Lake Louise, AB, 1991), NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., Kluwer Acad. Publ., Dordrecht, 1993.
- [5] M. Kolster, T. Nguyen Quang Do, and V. Fleckinger, *Twisted S -units, p -adic class number formulas, and the Lichtenbaum conjectures*, Duke Math. J. **84** (1996), 679–717.
- [6] T. Kubota and H.W. Leopoldt, *Eine p -adische Theorie der Zetawerte, I. Einführung der p -adischen Dirichletschen L -Funktionen*, J. reine angew. Math. **214/215** (1964), 328–339.
- [7] M. Kurihara, *Some remarks on conjectures about cyclotomic fields and K -groups of \mathbf{Z}* , Compositio Math. **81** (1992), 223–236.
- [8] C. Soulé, *K -théorie des anneaux d'entiers de corps de nombres et cohomologie étale*, Invent. Math. **55** (1979), 251–295.
- [9] H. Sumida-Takahashi, *The Iwasawa invariants and the higher K -groups associated to real quadratic fields*, Exp. Math. **14** (2005), 307–316.
- [10] H. Sumida-Takahashi, *Computation of the p -part of the ideal class group of certain real abelian fields*, Math. Comp. **76** (2007), 1059–1071.