



Supplementary Figure 1




Supplementary Figure 1: Analysis of monocyte subsets and lineage relationships. (a) Gating strategy for definition of MDP and cMoP populations in BM of *Cx3cr1^{GFP/+}* mice related to **Fig. 1a**. MDP was defined as CD11c⁻MHC-II⁻ cells from live CD45⁺Lin⁻CD11b⁻CD117⁺CD115⁺GFP⁺ population. For cMoP Ly6C⁺CD11c⁻MHC-II⁻ subset was gated from live CD45⁺Lin⁻CD11b⁻CD117⁺CD115⁺GFP⁺ population. (Lin: CD3, CD45R/B220, CD19, NK1.1, Ly6G, CD49b). **(b)** Gating strategy for identification of monocyte and GC (top) or MDP and cMoP (bottom) subpopulations in wild-type (*Cx3cr1^{+/+}*) mice.

Supplementary Figure 2




Supplementary Figure 2: Validation of gating strategy in *Nr4a1-GFP* mice and adoptive transfer experiments. (a) Flow cytometry plot depicting validation of CX₃CR1 based gating strategy (corresponding to gating in *Cx3cr1*^{GFP/+} mice, in **Fig. 1a**) precisely identifies Nr4a1-GFP⁺CD43⁺Ly6C^{lo/-} monocytes in bone marrow. (b) Flow cytometry plot depicting validation of CX₃CR1 independent gating strategy (corresponding to gating in wild type mice, in **Supplementary Fig 1b**) precisely defines Nr4a1-GFP⁺CD43⁺Ly6C^{lo/-} monocytes.

Supplementary Figure 3




Supplementary Figure 3: Ly6C^{hi} monocyte adoptive transfer experiments. (a) Sorting strategy for adoptive transfer of CD11b⁺GFP⁺Ly6C^{hi} monocytes. (b) Schematic illustration of adoptive transfer and multicolor flow cytometry analysis of Ly6C^{hi} monocytes from GFP⁺CD45.2⁺ mice into CD45.1⁺ recipients. (c) 4th and 5th rows of flow cytometry plots from **Fig. 1d** are shown. Donor CD45.2⁺CD11b⁺GFP⁺ cells are black. For comparison CD45.1⁺CD11b⁺Ly6C^{lo}/F4/80^{lo} recipient monocytes are depicted in blue (representative of two experiments).

Supplementary Figure 4




Supplementary Figure 4: Specificity and efficiency of Cre expression and Notch2 targeting. (a) Flow cytometry plot showing gradual expression of *LysM-eGFP* in myeloid cell subpopulations from cMoP to *Ly6C*^{lo} monocytes. (b) Flow cytometry plot showing YFP expression in bone marrow of *LysM*^{Cre}*Rosa*^{YFP} or control *Rosa*^{YFP} mice. *LysM*^{Cre} is strongly active in monocytes (*Ly6C*^{hi} and *Ly6C*^{lo}) and granulocytes but not in other cell populations. (c) Quantitative RT-PCR in sorted bone marrow cells from *GFP*⁺*Notch2*^{ΔMy} mice showing highest expression of *Cre* in N2⁺*Ly6C*^{hi} and subsequent reduction in *Ly6C*^{lo} monocytes (n=6). * *P* < 0.05, ** *P* < 0.01, *** *P* < 0.001; 1way ANOVA with Bonferroni's multiple comparison test. Error bars represent s.e.m. (d) *Notch2* expression in *Ly6C*^{hi} and *Ly6C*^{lo} monocyte subpopulations isolated from BM of *Notch2*^{ΔCD11c} mice (representative of two experiments). Littermate controls are shown for comparison.

Supplementary Figure 5




Supplementary Figure 5: Myeloid cell populations in *Notch2* deficient mice. (a) Monocyte subpopulation analysis based on CD115⁺ gating strategy reveals reduction of Ly6C^{lo} but not Ly6C^{hi} monocytes in *GFP⁺Notch2^{ΔMy}* mice. Data are pooled from three experiments (n=8/11). (b) Absolute or relative numbers of granulocytes in *GFP⁺Notch2^{ΔMy}* mice. Data are pooled from three experiments (n=8/10). (c) Relative frequency of monocytes by flow cytometry in *Notch2^{ΔMy}* with wild-type *Cx3cr1* locus. Data are pooled from three experiments (n=9/11). (d) Myeloid cell populations in *LysM^{Cre}Notch2^{+/+}* and *LysM^{+/+}Notch2^{+/+}* littermate control mice. Data are from three experiments (n=7/8). (e) Myeloid cell population analysis in *N1^{ΔMy}* mice (n=8/9). Data are pooled from three experiments. (f) Myeloid cell population analysis in *N1N2^{ΔMy}* mice. Data are pooled from two experiments (n=5/6). (a-f) * *P*<0.05, ** *P*<0.01, *** *P*<0.001; Student's t test. Error bars represent s.e.m.

Supplementary Figure 6




Supplementary Figure 6: Notch2 deficient BM Ly6C^{hi} monocytes show impaired conversion potential *in vivo*. (a, b) Flow cytometry (a) and corresponding fluorescence minus one controls (FMO ctrls) (b) 4 days after adoptive transfer of Ly6C^{hi} monocytes (corresponding to Fig. 4c) from control or Notch2 deficient CD45.2⁺GFP⁺ donors into CD45.1⁺ congenic recipients. (a) Transferred cells are shown in black and for comparison, recipient CD45.1⁺ (1st row), CD45.1⁺CD11b⁺ (2nd row), CD45.1⁺CD11b⁺Ly6C^{hi} (3rd row) cells or CD45.1⁺CD11b⁺Ly6C^{hi}F4/80^{lo} monocytes (4th and 5th rows) are depicted in blue.

Supplementary Figure 7




Supplementary Figure 7: Notch2 deficient peripheral Ly6C^{hi} monocytes show impaired conversion potential *in vivo*. (a) Flow cytometry plots 4 days after adoptive transfer of pooled splenic and peripheral blood Ly6C^{hi} monocytes from control or Notch2 deficient CD45.2⁺GFP⁺ donors into CD45.1⁺ congenic recipients. Transferred cells are shown in black and for comparison, recipient CD45.1⁺ (1st row), CD45.1⁺CD11b⁺ (2nd row), CD45.1⁺CD11b⁺Ly6C^{hi} (3rd row) cells or CD45.1⁺CD11b⁺Ly6C^{hi}F4/80^{lo} monocytes (4th to 5th rows) are depicted in blue (representative of two experiments).

Supplementary Figure 8




Supplementary Figure 8: Notch ligand DLL1 mediates monocyte conversion *in vitro*. (a) Gating strategy for definition and quantification of Ly6C^{lo} monocyte-like cells *in vitro*. Ly6C^{lo} monocyte-like cells (CD11b⁺GFP⁺Ly6C^{lo/-}CD11c^{lo}CD43⁺MHC-II^{lo/-}) are calculated as a percentage of live CD11b⁺GFP⁺ cells. (b) Flow cytometry plot showing expression of CD115 and upregulation of GFP on *in vitro* converted Ly6C^{lo} monocyte-like cells. *Ex vivo* isolated GFP⁺Ly6C^{hi} BM monocytes served as a staining control.

Supplementary Figure 9



Supplementary Figure 9: *In vivo* targeting of endothelial cells in mice. b(c) Specific β -galactosidase activity in capillaries and large vessels of peripheral muscle (top) and in aortic EC (en face preparations (bottom)) after treatment of *lacZ^{EC}* or control mice with tamoxifen; scale bar 100 μ m. Image is a representative of two (muscle) or three (aorta) experiments. (d) PCR for floxed or recombined locus of *Dll1* from heart (left), lung (middle) and peripheral muscle (right). Results are from one experiment representative of 4 experiments. (e) Quantitative RT-PCR analysis of *Dll1* expression in sorted ECs from control or *Dll1^{ΔEC}* mice. Pooled from three experiments (n=3). (f) Myeloid cell population analysis in *Dll1^{+/+}* or *CreERT2 Dll1^{+/+}* mice after tamoxifen treatment showing no influence of tamoxifen on Ly6C^{lo} monocyte development. Data are pooled from two experiments (n=6/7). (g) Cell population analysis in *Cdh5(PAC)-CreERT2 Dll1^{fl/fl}* mice showing no influence of *Cdh5(PAC)-CreERT2* expression on Ly6C^{lo} monocyte development. Data are pooled from three experiments (n=6/8). (e-g) * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$; Student's t test. Error bars represent s.e.m.

Supplementary Figure 10



Supplementary Figure 10: *In vivo* targeting of arterial endothelial cells in mice. (a) β -galactosidase staining demonstrates specific staining in central arteries of splenic follicles, arteries of peripheral muscle and aorta in *lacZ^{iaEC}* mice. Scale bars 100 μ m. Results are from one experiment representative of two independent experiments. (b) PCR for floxed or recombined *Dll1* locus ($\Delta Dll1$) from aortas (n=2).

Supplementary Table 1: Surface phenotype signatures for identification of distinct myeloid populations *in vivo*

Population	Phenotype
MDP	Lin ⁻ CD117 ⁺ CD11b ⁻ CD115 ⁺ CX ₃ CR1 ⁺ Ly6C ⁻ F4/80 ⁻ CD11c ⁻ MHC-II ⁻ or Lin ⁻ CD117 ⁺ CD11b ⁻ CD115 ⁺ Ly6C ⁻ F4/80 ⁻ CD11c ⁻ MHC-II ⁻
cMoP	Lin ⁻ CD117 ⁺ CD11b ⁻ CD115 ⁺ CX ₃ CR1 ⁺ Ly6C ^{hi} F4/80 ⁻ CD11c ⁻ MHC-II ⁻ or Lin ⁻ CD117 ⁺ CD11b ⁻ CD115 ⁺ Ly6C ^{hi} F4/80 ⁻ CD11c ⁻ MHC-II ⁻
Ly6C ^{hi}	Lin ⁻ CD117 ⁻ CD11b ⁺ CX ₃ CR1 ^{lo} Ly6C ^{hi} F4/80 ^{lo/-} CD11c ⁻ MHC-II ^{lo/-} CD43 ⁻ or Lin ⁻ CD117 ⁻ CD11b ⁺ Ly6C ^{hi} F4/80 ^{lo/-} CD11c ⁻ MHC-II ^{lo/-} CD43 ⁻
Ly6C ^{lo}	Lin ⁻ CD117 ⁻ CD11b ⁺ CX ₃ CR1 ^{hi} Ly6C ^{lo/-} F4/80 ^{lo} CD11c ^{lo} MHC-II ^{lo/-} CD43 ⁺ or Lin ⁻ CD117 ⁻ CD11b ⁺ Ly6C ^{lo/-} F4/80 ^{lo} CD11c ^{lo} MHC-II ^{lo/-} CD43 ⁺
GC	Lin ⁺ CD11b ⁺ CX ₃ CR1 ⁻ Ly6C ^{lo} or Lin ⁺ CD11b ⁺ Ly6C ^{lo}
Atypical cells (Ly6C ^{lo})	Lin ⁻ CD117 ⁻ CD11b ⁺ CX ₃ CR1 ^{hi} Ly6C ^{lo/-} F4/80 ^{lo} CD11c ⁻ MHC-II ^{hi} CD43 ⁻ or Lin ⁻ CD117 ⁻ CD11b ⁺ Ly6C ^{lo/-} F4/80 ^{lo} CD11c ⁻ MHC-II ^{hi} CD43 ⁻

Lin: CD3, CD45R/B220, CD19, NK1.1, Ly6G, CD49b

Supplementary Table 2: Mouse models used in the study

Abbreviations	Mouse description	Mouse background
CD45.1 ⁺	B6.SJL- <i>Ptprc</i> ^a <i>Pepec</i> ^b /BoyJ	B6
<i>GFP</i> ⁺ <i>ctrl</i>	<i>LysM</i> ^{+/+} <i>Notch2</i> ^{lox/lox} <i>Cx3cr1</i> ^{GFP/+}	B6
<i>GFP</i> ⁺ <i>N2</i> ^{ΔMy}	<i>LysM</i> ^{Cre} <i>Notch2</i> ^{lox/lox} <i>Cx3cr1</i> ^{GFP/+}	B6
<i>Ctrl</i>	<i>LysM</i> ^{+/+} <i>Notch2</i> ^{lox/lox}	B6
<i>N2</i> ^{ΔMy}	<i>LysM</i> ^{Cre} <i>Notch2</i> ^{lox/lox}	B6
<i>Ctrl</i>	<i>LysM</i> ^{+/+} <i>Notch1</i> ^{lox/lox}	B6
<i>N1</i> ^{ΔMy}	<i>LysM</i> ^{Cre} <i>Notch1</i> ^{lox/lox}	B6
<i>Ctrl</i>	<i>LysM</i> ^{+/+} <i>Notch1</i> ^{lox/lox} <i>Notch2</i> ^{lox/lox}	B6
<i>N1N2</i> ^{ΔMy}	<i>LysM</i> ^{Cre} <i>Notch1</i> ^{lox/lox} <i>Notch2</i> ^{lox/lox}	B6
<i>wt</i>	<i>Dll1</i> ^{+/+}	129
<i>Dll1</i> ^{+/lacZ}	<i>Dll1</i> ^{+/lacZ}	129
<i>Ctrl</i>	<i>Gt(ROSA)26Sor</i>	B6
<i>lacZ</i> ^{iEC}	<i>Cdh5(PAC)-CreERT2</i>	B6
<i>lacZ</i> ^{iaEC}	<i>Gt(ROSA)26Sor</i> <i>Bmx(PAC)-CreERT2</i>	B6
<i>Ctrl</i>	<i>Dll1</i> ^{lox/lox}	Mixed, B6;129
<i>Dll1</i> ^{iΔEC}	<i>Cdh5(PAC)-CreERT2 Dll1</i> ^{lox/lox}	Mixed, B6;129
<i>Dll1</i> ^{iΔaEC}	<i>Bmx(PAC)-CreERT2 Dll1</i> ^{lox/lox}	Mixed, B6;129
<i>Ctrl</i>	<i>Dll4</i> ^{lox/lox}	Mixed, B6;CD1
<i>Dll4</i> ^{iΔEC}	<i>Cdh5(PAC)-CreERT2 Dll4</i> ^{lox/lox}	Mixed, B6;CD1
<i>Dll1</i> ^{+/+}	<i>Dll1</i> ^{+/+}	Mixed, B6;129
<i>CreERT2 Dll1</i> ^{+/+}	<i>Cdh5(PAC)-CreERT2 Dll1</i> ^{+/+}	Mixed, B6;129
<i>Ctrl</i>	<i>LysM</i> ^{+/+} <i>Notch2</i> ^{+/+}	B6
<i>LysM</i> ^{Cre}	<i>LysM</i> ^{Cre} <i>Notch2</i> ^{+/+}	B6
<i>Nr4a1-GFP</i>	<i>Nr4a1-EGFP/Cre</i>	B6
<i>LysM-eGFP</i>	<i>LysM-EGFP</i>	B6
<i>Rosa</i> ^{YFP}	<i>LysM</i> ^{+/+} <i>Gt(ROSA)YFP26Sor</i>	B6
<i>LysM</i> ^{Cre} ; <i>Rosa</i> ^{YFP}	<i>LysM</i> ^{Cre} <i>Gt(ROSA)YFP26Sor</i>	B6
<i>GFP</i> ⁺ <i>ctrl</i>	<i>CD11c</i> ^{Cre} <i>Notch1</i> ^{lox/+} <i>Notch2</i> ^{lox/+} <i>Cx3cr1</i> ^{GFP/+}	B6
<i>GFP</i> ⁺ <i>N1N2</i> ^{ΔCD11c}	<i>CD11c</i> ^{Cre} <i>Notch1</i> ^{lox/lox} <i>Notch2</i> ^{lox/lox} <i>Cx3cr1</i> ^{GFP/+}	B6
<i>Ctrl</i>	<i>CD11c</i> ^{+/+} <i>Notch2</i> ^{lox/lox}	B6
<i>N2</i> ^{ΔCD11c}	<i>CD11c</i> ^{Cre} <i>Notch2</i> ^{lox/lox}	B6

Supplementary Table 3: Antibodies and fluorescence dyes for flow cytometry and immunofluorescence used in the study

Antibody/dye	Clone	Dilution	Company
Anti-CD3ε	145-2C11	1:100	BioLegend
Anti-CD49b	DX5	1:400	BioLegend
Anti-CD45R/B220	RA3-6B2	1:400	BioLegend
Anti-Ly6G	1A8	1:400	BioLegend
Anti-CD19	1D3	1:400	BD Pharmingen
Anti-NK1.1	PK136	1:400	BioLegend
Anti-CD117	2B8	1:100	BioLegend
Anti-CD115	AFS98	1:100	BioLegend
Anti-CD11b	M1/70	1:400	BioLegend
Anti-Ly6C	HK1.4	1:2800	BioLegend
Anti-F4/80	BM8	1:100	BioLegend
Anti-CD11c	N418	1:400	BioLegend
Anti-I-A/I-E	M5/114.15.2	1:400	BioLegend
Anti-CD43	S7	1:400	BD Pharmingen
Anti-CD45	30-F11	1:200	BioLegend
Anti-CD45.1	A20	1:100	BioLegend
Anti-CD45.2	104	1:200	BioLegend
Anti-CD144	11D4.1	1:100	BD Pharmingen
Anti-CX ₃ CR1	SA011F11	1:200	BioLegend
Anti-CD11a	M17/4	1:200	BioLegend
Anti-CCR2	475301	1:100	R&D
Anti-Notch2	HMN2-35	1:100	BioLegend
Anti-CD16/CD32	93	1:200	BioLegend
Anti-CD31	Mec13.3	1:400	BD Pharmingen
Streptavidin PE-Dazzle594		1:400	BioLegend
Annexin V		1:50	BioLegend
7AAD		1:100	BioLegend
Propidium Iodide		1:12000	Sigma
Anti-GFP		1:300	Acris
Anti-DLL1	HMD1-3	1:100	BioLegend
Streptavidin Cy3		1:400	BioLegend
Anti-rabbit IgG-FITC		1:200	Jackson ImmunoResearch
DAPI		1:5000	Invitrogen

Supplementary Table 4: Primers for QRT-PCR

Gene	Primers
<i>Notch2</i>	Forward: AGTGTCAGAGGCCAGCAAGAAGAA Reverse: TGATTGTCGTCCATCAGAGCACCA
<i>Notch1</i>	Forward: TGGAGGTCTCAGTGGCTATAA Reverse: ATTCTGGCATGGGTTAGAAAGA
<i>Hey2</i>	Forward: TGAAGCGCCCTTGTGAGGAA Reverse: TTGTAGCGTGCCCAGGGTAA
<i>Hes1</i>	Forward: CCGGACAAACCAAAGACGGC Reverse: GGAATGCCGGGAGCTATCTTTCT
<i>Dll1</i>	Forward: TCCGATTCCCCTTCGGCTTC Reverse: TGGGTTTTCTGTTGCGAGGT
<i>Nr4a1</i>	Forward: AGCTTGGGTGTTGATGTTCC Reverse: AATGCGATTCTGCAGCTCTT
<i>Pparg</i>	Forward: AGGGCGATCTTGACAGGAA Reverse: CACCTCTTTGCTCTGCTCCT
<i>Pou2f2</i>	Forward: TGCACATGGAGAAGGAAGTG Reverse: AGCTTGGGACAATGGTAAGG
<i>Sfn5</i>	Forward: AGGTCGAACGATTCTGCTGT Reverse: TCTGAGGGAAACTGGAAAGG
<i>Ccr2</i>	Forward: CCTTGGGAATGAGTAACTGTGTGAT Reverse: ATGGAGAGATACCTTCGGAACTTCT
<i>Cx3cr1</i>	Forward: GCAGAAGTTCCTTCCCATC Reverse: GGACAGGAAGATGGTTCCAA