

## CASE REPORT

# An Adolescent Athlete with Low Back Pain Associated with Spina Bifida Occulta at the Thoracolumbar Junction : A Case Report

Tsuyoshi Goto<sup>1</sup>, Toshinori Sakai<sup>2</sup>, Nori Sato<sup>1</sup>, Shinsuke Katoh<sup>1</sup>, and Koichi Sairyo<sup>2</sup>

<sup>1</sup>Department of Rehabilitation Medicine, Tokushima University Hospital, Tokushima, Japan, <sup>2</sup>Department of Orthopedics, Institute of Biomedical Sciences, Tokushima University Graduate School, Tokushima, Japan

**Abstract : Introduction :** Spina bifida occulta (SBO) is a common malformation of lamina of the spine, most commonly occurring in the sacrum or lower lumbar spine. Athletes with anatomical abnormalities such as SBO may be vulnerable to discoordination of muscles and/or skeletal structures. **Case Report :** We present a case of a 15-year-old male adolescent who presented with persistent low back pain (LBP) associated with SBO at T11-L1 that had been overlooked for several years. He participated in karate as a top-level competitor at the national level, but the LBP interfered with his play. Dynamic X-ray films showed limited range of motion (ROM) in his lower lumbar spine (L3-S1). Instead, the thoracolumbar junction compensated for the lack of ROM in the lumbar spine, enabling trunk flexion and extension. To restore normal coordination, he was treated with a 2-week rehabilitation program including flexibility and core conditioning for the trunk to improve coordination of spinal movement. Two weeks later, his LBP resolved. **Conclusion :** These results suggest that evaluation of the coordination of the muscles and skeletal structures has an important role in the treatment of cases with an anatomical abnormality such as SBO. *J. Med. Invest.* 66 : 199-200, February, 2019

**Keywords :** adolescent athlete, low back pain, spina bifida occulta, thoracolumbar junction

## INTRODUCTION

Spina bifida occulta (SBO) is a common malformation of lamina of the spine, most commonly occurring in the sacrum or lower lumbar spine (1). It is rarely found or may be overlooked at other spinal levels. In SBO, the spinal column is thought to be affected by the lack of continuity of the posterior elements and/or attachments of associated muscles. Athletes with anatomical abnormalities such as SBO may be vulnerable to discoordination of muscles and/or skeletal structures. In this report, we present a case of an adolescent with chronic low back pain (LBP) associated with SBO at T11-L1, which had been overlooked for several years.

## CASE REPORT


A 15-year-old male adolescent presented with a 3-year history of LBP that was induced when his trunk was rotated to the left and bent to forward. He participated in karate as a top-level competitor at the national level but the LBP interfered with his play. Several doctors he had seen had diagnosed this as LBP of unknown origin.

On radiological examinations including X-ray films and computed tomography (CT) scans, there was a lack of spinous processes at T11-L1, indicating spina bifida occulta (SBO) (Figure 1a, 1b). MRI revealed no obvious abnormal findings except the SBO. Dynamic X-ray films showed limited range of motion (ROM) in his lower lumbar spine (L3-S1). Instead, the thoracolumbar junction

compensated for the lack of ROM in the lumbar spine, enabling trunk flexion and extension (Figure 1b).

We believed that the discoordination of muscles and skeletal structures derived from the SBO at T11-L1 and was the main cause of his LBP. To restore normal coordination, he was treated with a 2-week rehabilitation program including flexibility and core conditioning for the trunk.

We evaluated his spinal motion using a SpinalMouse® (Idiag, Volkswill, Switzerland). The SpinalMouse® was used by having



**Figure 1** (a) Three-dimensional reconstructed computed tomography images showing spina bifida occulta at T11-L1. (b) Dynamic plain X-ray films of the whole spine showing decreased range of motion in the lower lumbar spine.

Received for publication August 2, 2018 ; accepted October 25, 2018.

Address correspondence and reprint requests to Toshinori Sakai, MD, PhD, Associate Professor, Department of Orthopedics, Institute of Biomedical Sciences, Tokushima University Graduate School, 3-18-15 Kuramoto-cho, Tokushima 770-8503, Japan and Fax : +81-88-633-0178

the patient roll paravertebrally from the 7th cervical (C7) to the 3rd sacral (S3) vertebra. Using the SpinalMouse<sup>®</sup>, we also measured each intersegmental angle while the patient was in the following positions : a neutral crawling on hands and knees position (NCP), a maximum extension position (MEP), and a maximum flexion position (MFP) to evaluate thoracic, lumbar, and total spinal mobility (Figure 2). In addition, motion analyses were performed using the Vicon system (Vicon Motion Systems, Oxford, United Kingdom) before and 2 weeks after beginning the rehabilitation program.

Two weeks after the patient began the rehabilitation program, thoracic and lumbar mobility improved from 77° to 86° and 17° to 31°, respectively, on the SpinalMouse<sup>®</sup>. The thoracic extension angle and spine extension angle improved from 4.6° to 2.9° and 29.0° to 17.5°, respectively, according to the Vicon system. His LBP improved from 56 to 21/100 on a visual analogue scale.

He has continued with this conditioning for 2 years since the first visit to our clinic. He has been able to continue participating in karate as a top-level competitor at the national level, with no LBP.

## DISCUSSION

Spina bifida occulta (SBO), which is frequently seen in the lumbosacral area, is considered a developmental failure of osseous union between the two halves of the posterior arch. In this report, we presented a case with SBO in the thoracolumbar junction. To our knowledge, no report has described SBO in the thoracolumbar junction, though the probable incidence is regarded as < 5% (2). The thoracolumbar spine is the focal point of stress during various spinal movements. Specifically, the central point of this girdling

structure is the thoracolumbar fascia, which is a complex of aponeurotic and fascial planes that forms the retinaculum around the paraspinal muscles of the lower back and sacral region (3). This complex is continuous with the paraspinal fascia in the cervical, thoracic, and lumbo-pelvic regions, eventually fusing to the cranial base and iliac crest. Based on these anatomical features, a loss of normal structure in the thoracolumbar spine may cause discoordination of motion in this region and related functions, which results in LBP. Proctor, *et al.* reported a case series on thoracolumbar syndrome as a cause of LBP (4).

In this case, we hypothesized that discoordination of the muscles and skeletal structures caused by the SBO at T11-L1 were a main factor in his LBP. Generally, the coordination pattern between adjacent joint angles reflects motor control of the organizing adjacent structures. Accordingly, in this case, immediately after the spinal mobility measured by SpinalMouse<sup>®</sup> improved by the rehabilitation program, his LBP has disappeared. This result suggests that the coordination of the muscles and skeletal structures that was compromised, due to the SBO at T11-L1, had recovered.

## CONCLUSION

These results suggest that evaluation of the coordination of the muscles and skeletal structures has an important role in the treatment of cases with an anatomical abnormality such as SBO.

## DISCLOSURE

All authors confirm that there are no conflicts of interest with regard to persons or organizations that could have influenced our study findings.

## REFERENCES

1. Sakai T, Goda Y, Tezuka F, Abe M, Yamashita K, Takata Y, Higashino K, Nagamachi A, Sairyo K : Clinical features of patients with pars defects identified in adulthood. *Eur J Orthop Surg Traumatol* 26 : 259-262, 2016
2. Yochum TR, Rowe LJ : *Essentials of Skeletal Radiology*. 3rd ed. Lippincott Williams & Wilkins : Philadelphia, 2005, pp.309-310
3. Schuenke MD, Vleeming A, Van Hoof T, Willard FH : A description of the lumbar interfascial triangle and its relation with the lateral raphe : anatomical constituents of load transfer through the lateral margin of the thoracolumbar fascia. *J Anat* 221 : 568-576, 2012
4. Proctor D, Dupuis P, Cassidy JD : Thoracolumbar syndrome as a cause of low-back pain : A report of two cases. *J Can Chiropr Assoc* 29 : 71-73, 1985




Figure 2 SpinalMouse<sup>®</sup> images of the thoracolumbar spine showing a change in spinal movement from pre- to post-intervention. An increase in the lumbar flexion angle was especially noteworthy (circle).