

The novel inhibitor PRI-724 for Wnt/ β -catenin/CBP signaling ameliorates bleomycin-induced pulmonary fibrosis in mice

| | |
|-------------------------------|---|
| Journal: | <i>Experimental Lung Research</i> |
| Manuscript ID | UELR-2019-0012.R2 |
| Manuscript Type: | Original Paper |
| Date Submitted by the Author: | n/a |
| Complete List of Authors: | Okazaki, Hiroyasu; Graduate School of Biomedical Sciences, Tokushima University, Department of Respiratory Medicine and Rheumatology Sato, Seidai; Graduate School of Biomedical Sciences Tokushima University, Department of Respiratory Medicine and Rheumatology Koyama, Kazuya; Graduate School of Biomedical Sciences, Tokushima University, Department of Respiratory Medicine and Rheumatology Morizumi, Shun; Graduate School of Biomedical Sciences, Tokushima University, Department of Respiratory Medicine and Rheumatology Abe, Shuichi; Graduate School of Biomedical Sciences, Tokushima University, Department of Respiratory Medicine and Rheumatology Azuma, Momoyo; Graduate School of Biomedical Sciences, Tokushima University, Department of Respiratory Medicine and Rheumatology Chen, Yajuan; Graduate School of Biomedical Sciences, Tokushima University, Department of Respiratory Medicine and Rheumatology Goto, Hisatsugu; Graduate School of Biomedical Sciences, Tokushima University, Department of Respiratory Medicine and Rheumatology Aono, Yoshinori; Graduate School of Biomedical Sciences, Tokushima University, Department of Respiratory Medicine and Rheumatology; National Hospital Organization Higashi Tokushima Medical Center Ogawa, Hirohisa; Tokushima University Graduate School, Department of Pathology and Laboratory Medicine Kagawa, Kozo; Graduate School of Biomedical Sciences, Tokushima University, Department of Respiratory Medicine and Rheumatology Nishimura, Haruka; Graduate School of Biomedical Sciences, Tokushima University, Department of Respiratory Medicine and Rheumatology Kawano, Hiroshi; Graduate School of Biomedical Sciences, Tokushima University, Department of Respiratory Medicine and Rheumatology Toyoda, Yuko; Graduate School of Biomedical Sciences, Tokushima University, Department of Respiratory Medicine and Rheumatology Uehara, Hisanori; Tokushima University Graduate School, Department of Pathology and Laboratory Medicine |

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

| | |
|-----------|--|
| | Kouji, Hiroyuki; Oita University, Faculty of Medicine Nishioka, Yasuhiko; Graduate School of Biomedical Sciences, Tokushima University, Department of Respiratory Medicine and Rheumatology |
| Keywords: | Wnt/ β -catenin/CBP signaling, Idiopathic pulmonary fibrosis, Alveolar macrophage, PRI-724 |
| | |



1
2
3
4
5
6
7 The novel inhibitor PRI-724 for Wnt/ β -catenin/CBP signaling ameliorates
8
9
10 bleomycin-induced pulmonary fibrosis in mice
11
12
13
14
15

16 Hiroyasu Okazaki¹, Seidai Sato¹, Kazuya Koyama¹, Shun Morizumi¹, Shuichi Abe¹,
17
18 Momoyo Azuma¹, Yajuan Chen¹, Hisatsugu Goto¹, Yoshinori Aono^{1,2}, Hirohisa Ogawa³,
19
20
21
22 Kozo Kagawa¹, Haruka Nishimura¹, Hiroshi Kawano¹, Yuko Toyoda¹, Hisanori Uehara³,
23
24
25
26
27 Hiroyuki Kouji^{4,5}, Yasuhiko Nishioka¹
28
29
30
31
32

33 ¹Department of Respiratory Medicine and Rheumatology, Graduate School of
34
35
36 Biomedical Sciences, Tokushima University, Tokushima, Japan
37
38

39
40 ²National Hospital Organization Higashi Tokushima Medical Center, Tokushima, Japan
41
42

43 ³Department of Pathology and Laboratory Medicine, Tokushima University Graduate
44
45
46 School, Tokushima, Japan
47
48

49
50 ⁴PRISM BioLab Co., Ltd., 4259-3, Nagatsuda-cho, Midori-ku, Yokohama 226-8510,
51
52
53 Japan
54
55

56 ⁵Oita University, Faculty of Medicine, 1-1, Idaigaoka, Hasama-machi, Yufu-city, Oita
57
58
59
60

1
2
3
4
5
6
7 879-5593, Japan
8
9
10
11
12

13 Address correspondence and reprint requests to: Yasuhiko Nishioka, M.D., Ph.D.,
14

15
16 Department of Respiratory Medicine and Rheumatology, Graduate School of Biomedical
17

18
19 Sciences, Tokushima University, 3-18-15 Kuramoto-cho, Tokushima 770-8503, Japan.
20
21

22
23 Phone: +81-88-633-7127; Fax: +81-88-633-2134
24
25

26
27 E-mail: yasuhiko@tokushima-u.ac.jp
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Purpose/Aim of the Study: Wnt/ β -catenin signaling was reported to be activated in pulmonary fibrosis, and was focused on as a target for antifibrotic therapy. However, the mechanism how the inhibition of Wnt/ β -catenin signaling ameliorate pulmonary fibrosis has not been fully elucidated. The purpose of this study is to explore the target cells of Wnt/ β -catenin inhibition in pulmonary fibrosis and to examine the antifibrotic effect of the novel inhibitor PRI-724 specifically disrupting the interaction of β -catenin and CBP.

Materials and Methods: The effect of C-82, an active metabolite of PRI-724, on the expression of TGF- β 1 and α -smooth muscle actin (SMA) was examined on fibroblasts and macrophages. We also examined the effects of PRI-724 in mouse model of bleomycin-induced pulmonary fibrosis.

Results: The activation and increased accumulation of β -catenin in the canonical pathway were detected in lung fibroblasts as well as macrophages stimulated by Wnt3a using Western blotting. Treatment with C-82 reduced CBP protein and increased p300 protein binding to β -catenin in the nucleus of lung fibroblasts. In addition, C-82 inhibited the expression of SMA in lung fibroblasts treated with TGF- β , indicating the inhibition of

1
2
3
4
5
6 myofibroblast differentiation. In the fibrotic lungs induced by bleomycin, β -catenin was
7
8
9
10 stained strongly in macrophages, but the staining of β -catenin in alveolar epithelial cells
11
12
13 and fibroblasts was weak. The administration of PRI-724 ameliorated pulmonary fibrosis
14
15
16 induced by bleomycin in mice when administered with a late, but not an early, **treatment**
17
18
19 **schedule**. Analysis of bronchoalveolar fluid (BALF) showed a decreased number of
20
21
22 alveolar macrophages. In addition, the level of TGF- β 1 in BALF was decreased in mice
23
24
25 treated with PRI-724. C-82 also inhibited the production of TGF- β 1 by alveolar
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conclusions: These results suggest that the β -catenin/CBP inhibitor PRI-724 is a potent antifibrotic agent that acts by modulating the activity of macrophages in the lungs.

Key Words: Wnt/ β -catenin/CBP signaling, Idiopathic pulmonary fibrosis, Alveolar macrophage, PRI-724.

INTRODUCTION

1
2
3
4
5
6
7 Idiopathic pulmonary fibrosis (IPF) is a chronic devastating lung disease, characterized
8
9
10 by the proliferation of fibroblasts and deposition of extracellular matrix in the lungs (1,
11
12
13 2). The prognosis associated with IPF is poor, and the median survival time after
14
15
16 diagnosis is approximately 2 to 4 years (1). Although there are some antifibrotic agents
17
18
19 available for the treatment of IPF (3), the efficacy is not yet sufficient for patients because
20
21
22 the advantage is limited in terms of ameliorating the decline in the lung function, but not
23
24
25 the survival benefit or recovery of injured lungs. Thus, the development of novel drugs
26
27
28 with superior effects is strongly expected.
29
30
31
32

33 The pathogenesis of IPF is based on the repeated injury of alveolar epithelial cells
34
35
36 (AECs) and subsequent activation of mesenchymal cells with the formation of
37
38
39 fibroblastic and myofibroblastic foci (2, 4). The Wnt/ β -catenin pathway is well-known as
40
41
42 essential signaling during lung development and homeostasis (5). Activated Wnt/ β -
43
44
45 catenin signaling is associated with fibrosis in the liver, heart, kidney, skin, and other
46
47
48 organs including the lung (5, 6). In the lungs of IPF patients, Wnt/ β -catenin signaling is
49
50
51 activated aberrantly in AECs and mesenchymal cells (fibroblasts/myofibroblasts) (7-9).
52
53
54 Experimentally, the signaling is known to be induced in a murine model of lung fibrosis
55
56
57
58
59
60

1
2
3
4
5
6
7 with the intratracheal administration of bleomycin (10-12).
8
9

10 However, the role of Wnt/ β -catenin signaling in pulmonary fibrosis is complicated due
11
12 to the pleiotropic action in many types of lung cells. Increased β -catenin in AECs is
13
14 related to the hyperplasia and regeneration in IPF (7, 9, 11). It was reported that the
15
16 Wnt/ β -catenin pathway plays a role in the survival, migration, and proliferation of AECs
17
18 (9-12). The specific deletion of β -catenin in AECs resulted in the enhanced apoptosis of
19
20 AECs, leading to an increased number of fibroblasts and delayed recovery from epithelial
21
22 injury in mice (12). However, the activation of Wnt/ β -catenin signaling was also reported
23
24 to be related to epithelial mesenchymal transition (EMT), indicating, on the contrary, the
25
26 pro-fibrotic activity of Wnt/ β -catenin (11). In addition, lung fibroblast cell lines are
27
28 activated via Wnt/ β -catenin signaling in migration, proliferation, and collagen production
29
30 (8, 9, 11, 13). These results suggest that the inhibition and/or modulation of Wnt/ β -
31
32 catenin signaling may be applicable as a therapeutic strategy for pulmonary fibrosis. In
33
34 fact, several reports demonstrated that the inhibition of Wnt/ β -catenin signaling with
35
36 inhibitors (14-18) or inhibitory protein, such as Dickkopf-1 (18), ameliorated pulmonary
37
38 fibrosis in a bleomycin model.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7 However, more recently, macrophages were also reported to be novel cells that are
8
9
10 involved in the regulation of pulmonary fibrosis via the Wnt/ β -catenin signaling pathway
11
12
13 (19). In the present study, we performed cell-based analysis of the distribution and
14
15
16 activation of β -catenin in fibrotic lungs, and demonstrated the antifibrotic effects and
17
18
19 mode of action of the second-generation inhibitor PRI-724 specific for CBP/ β -catenin,
20
21
22
23 which has been used in clinical trials to treat patients with malignancy, in a bleomycin-
24
25
26 induced pulmonary fibrosis model in mice.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

METHOD

Reagents.

PRI-724 and the active metabolite C-82 were from PRISM BioLab, Co., Ltd (Kanagawa, Japan). Recombinant Mouse Wnt3a Protein (R&D Systems, Minneapolis, MN, USA), recombinant TGF- β 1 protein (R&D Systems), and Proteinase K (Sigma-Aldrich, Saint Louis, MO, USA) were also purchased.

Animals and Treatments.

Eight-week-old C57BL/6 male mice were purchased from Charles River Japan, Inc. (Yokohama, Japan). Mice were maintained in the animal facility of the University of Tokushima under specific pathogen-free conditions according to the guidelines of our university (20). The present study was approved by the Institutional Animal Care and Use Committee of Tokushima University (Permission Number: 14099). Bleomycin sulfate (mixture) (Tokyo Chemical Industry Co., Tokyo, Japan) was dissolved in sterile saline and injected via oropharyngeal aspiration at a dose of 1.25 mg/kg of bleomycin in a total volume of 50 μ L of saline. During the injection process, mice were anesthetized with

1
2
3
4
5
6
7 2.5% isoflurane delivered in a box. Mice were suspended vertically on a stand for
8
9
10 oropharyngeal aspiration. Osmotic minipumps (Alzet Osmotic Pumps, DURECT
11
12 Corporation, CA, USA) containing PBS or 1 or 10 mg/kg/day of PRI-724 were implanted
13
14
15
16 subcutaneously.
17
18
19
20
21
22

23 **Cell Lines.**

24
25
26 Murine lung fibroblasts were generated from the lungs of C57BL/6 mice as reported
27
28 previously (20, 21). These fibroblasts were used at 5 to 10 passages. The human lung
29
30 fibroblast cell line MRC5 and macrophage cell line RAW257.6 were obtained from DS
31
32 PHARMA BIOMEDICAL (Osaka, Japan) and the American Type Culture Collection,
33
34 ATCC (Manassas, VA, USA), respectively. These cells were cultured with RPMI1640
35
36 medium supplemented with 10% fetal bovine serum (FBS) (GIBCO BRL, Rockville, MD,
37
38
39
40
41
42
43
44
45
46
47 USA).

53 **Bronchoalveolar Lavage.**

54
55
56 Bronchoalveolar Lavage (BAL) was performed with saline (1 mL) using a soft cannula
57
58
59
60

1
2
3
4
5
6 on mice treated with bleomycin (20). BAL fluid (BALF) was collected on days 28 and
7
8
9
10 35 after bleomycin injection, and centrifuged at 1,000 rpm for 10 minutes. The
11
12
13 supernatant was used for ELISA to measure the TGF- β 1 level. After counting the cell
14
15
16 number in the BALF, cells were cytopun onto glass slides and stained with Diff-Quick
17
18
19 (Baxter, Miami, FL, USA) for cell classification.
20
21
22
23
24
25
26

27 **Alveolar Macrophage Isolation.**

28
29
30 BAL was conducted for mice treated by bleomycin 7 days after administration. The cells
31
32
33 were plated in wells in 1.0% FBS RPMI1640 medium. One hour later, the wells were
34
35
36 washed with PBS three times and the adherent cells were used as alveolar macrophages
37
38
39 (AMs). AMs (1×10^6 /mL) were cultured on the well in 1.0% FBS RPMI1640 and the
40
41
42 supernatants were collected after 24 hours-treatment with 1.0 μ M of C-82 or DMSO.
43
44
45
46 Cells were collected for real-time PCR to examine the mRNA level.
47
48
49
50
51
52

53 **Hydroxyproline Assay.**

54
55
56 Quantitative examination of the collagen content was performed using the
57
58
59
60

1
2
3
4
5
6 hydroxyproline assay kit (BioVison, Milpitas, CA, USA) as previously described (22).
7
8

9
10 The left lobes or total lungs were used for this assay to allow comparison.
11
12
13
14
15

16 **Western Blot Analysis.**

17
18
19
20 Serum-starved lung fibroblast B6 cells were harvested in RPMI1640 with 0.1% FBS and
21
22
23 1% penicillin-streptomycin. Thirty micrograms of protein per sample was loaded onto
24
25
26 8% tris-glycine SDS polyacrylamide gels (Invitrogen) for electrophoresis and then
27
28
29 transferred onto polyvinylidene difluoride (PVDF) membranes (0.2 μ M, ATTO).
30
31
32
33 Membranes were then incubated with Blocking One (Nakalai Tesque Inc, Japan) for 1
34
35
36 hour at room temperature and then incubated with the appropriate primary antibody
37
38
39 overnight. Secondary antibodies and an ECL kit (GE) were applied for generating
40
41
42 chemiluminescent signals. Primary antibodies used for Western analysis included: anti-
43
44
45 α -SMA (Sigma-Aldrich, St. Louis, MO, USA), **anti-non-phospho (Active) β -catenin**
46
47
48 **antibody (Ser33/37/Thr41) (D13A1) (Cell Signaling Technology, Danvers, MA, USA),**
49
50
51
52
53 anti-phospho- β -Catenin (Y654) antibody (St John's Laboratory, London, United
54
55
56 Kingdom), anti-GAPDH (6C5) (Santa Cruz Biotechnology, Santa Cruz, CA, USA), anti-
57
58
59
60

1
2
3
4
5
6 LAMIN A/C (Cell Signaling Technology, Danvers, MA, USA), anti-p300 (C-20) (Santa
7 Cruz Biotechnology), and anti-CBP (A-22) (Santa Cruz Biotechnology) antibodies.
8
9
10
11
12
13 Normal rabbit IgG (Millipore) and Protein G-Sepharose 4 Fast flow (GE Healthcare,
14 Buckinghamshire, UK) were used. Densitometric analysis was performed using National
15
16
17 Institutes of Health (NIH) Image J software.
18
19
20
21
22
23
24
25
26

27 **Cytoplasmic and Nuclear Protein Extraction.**

28
29
30 Proteins from cytoplasmic and nuclear compartments were separated using NE-PER
31
32
33 Nuclear and Cytoplasmic Extraction Reagents (Thermo Fisher Scientific). Briefly,
34
35
36 fibroblast and macrophage cells were harvested with trypsin-EDTA and then washed
37
38
39
40 twice with PBS. Then, cells were centrifuged at $500 \times g$ for 5 minutes and supernatants
41
42
43 were removed. Ice-cold CER-I and CER-II solutions were added according to the
44
45
46 manufacturer's instructions to separate the cytoplasmic from the nuclear compartment
47
48
49
50 proteins. Western blot for GAPDH and LAMIN A/C were used to ensure no
51
52
53 contamination in each part of the extracts.
54
55
56
57

58 **Histopathology.**

1
2
3
4
5
6
7 The right lungs were fixed in 10% buffered formalin and embedded in paraffin. Sections
8
9
10 (3-4 μm) were stained with hematoxylin and eosin. Azan staining was also performed.
11
12
13
14
15

16 **TGF- β 1 ELISA.**

17
18
19
20 The concentrations of TGF- β 1 in BALF and the culture supernatant of macrophages were
21
22
23 determined using the TGF- β 1 ELISA kit according to the manufacturer's instructions
24
25
26 (R&D Systems). To determine the amount of total TGF- β 1 in the conditioned media,
27
28
29 samples were pretreated with 1 M HCl for 15 min at room temperature prior to
30
31
32 neutralization 1 M NaOH, as suggested by the manufacturer.
33
34
35
36
37
38
39

40 **Immunofluorescence Staining.**

41
42
43 Paraffin-embedded lung-section slides were de-paraffinized, rehydrated, and subjected to
44
45
46 antigen retrieval using proteinase K (20 $\mu\text{g}/\text{mL}$) for 25 minutes in 37°C before
47
48
49 immunostaining (23). Non-specific binding was blocked using a solution of 5% BSA with
50
51
52 0.3% Triton X-100 in TBS. The lung sections were incubated with primary antibodies to
53
54
55 **phospho-Y654- β -catenin**, nonphospho- β -catenin, CD68, pro-SPC, α -SMA, and Sca-1 in
56
57
58
59
60

1
2
3
4
5
6
7 1% BSA with 0.3% Triton X-100 in TBS. The sections were then washed and incubated
8
9
10 with Alexa Fluor-594 goat anti-rabbit IgG secondary antibody, and Alexa Fluor-488 anti-
11
12 rat IgG secondary antibody and nuclei were counterstained with DAPI. **Fluorescence**
13
14 analysis was performed using an Olympus inverted fluorescence microscope. Primary
15
16 antibodies used included: **anti-Active- β -catenin (anti-ABC) (Millipore, Massachusetts,**
17
18 **USA), anti-Sca-1/Ly6A/E (D7) (FITC) (Abcam, Cambridge, MA, USA), anti-pro-SPC**
19
20 **(Abcam), anti-CD68 (Abdserotec, Oxford, UK), and anti- α -SMA (1A4) (Biotin)**
21
22 **(Abcam) antibodies. Other reagents such as Streptavidin, Alexa Fluor 488 conjugate**
23
24 **(Thermo Fisher Scientific, Waltham, MA, USA) and APEX Alexa Fluor 488 antibody**
25
26 **labeling kit (Thermo Fisher Scientific) were also used.**
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 **Quantitative Real-time PCR Analysis.**

44
45
46 Total mRNA was extracted from cells using TRIzol reagent and cDNA was synthesized
47
48
49 from mRNA using the Prime Script RT-PCR synthesis kit (High Capacity RNA-to-cDNA
50
51
52 kit, Applied Biosystems) following the manufacturer's instructions. RT-PCR was
53
54
55 performed using the standard protocol. Primers detecting TGF- β 1 and GAPDH were
56
57
58
59
60

1
2
3
4
5
6
7 used:

8
9
10 TGF- β 1

11
12
13 For: CAACAATTCCTGGCGTTACCTTGG

14
15
16 Rev: GAAAGCCCTGTATTCCGTCTCCTT

17
18
19 GAPDH

20
21
22 For: CAACTACATGGTCTACATGTTC

23
24
25 Rev: CGCCAGTAGACTCCACGAC
26
27
28
29
30
31
32

33 **Statistical Analysis.**

34
35
36 Statistical analysis was performed with GraphPad Prism Ver. 5.01 (GraphPad Software
37
38
39 Inc.). Dates are expressed as the mean \pm standard error (SE). Differences in measured
40
41
42 variables between experimental and control groups were assessed using Student's t-test.
43
44
45

46 Values of $P < 0.05$ were considered significant.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RESULTS

Activation of β -catenin in lung fibroblasts and macrophages.

We first demonstrated the activation of β -catenin in lung fibroblasts and macrophages with the primary culture of lung fibroblasts called B6 cells and macrophage cell line RAW257.6. The activation of β -catenin was evaluated by the nuclear accumulation with the anti-active (nonphospho) form of β -catenin dephosphorylated on Ser37 or Thr41 and phospho (p)-Y654- β -catenin antibodies by Western blotting. In lung fibroblast B6 cells, an increased accumulation of nonphospho- β -catenin was detected in the nucleus and the cytoplasm one hour after Wnt3a stimulation (Figure 1A, Supplementary Figure 1). Similarly, in macrophage RAW257.6 cells, the accumulation of both nonphospho- and pY654- β -catenin was demonstrated 12 hours after Wnt3a stimulation (Figure 1B, Supplementary Figure 2, 3). These results suggest that the activation of β -catenin was observed not only in fibroblasts in addition to alveolar epithelial cells, but also in macrophages.

C-82, an active metabolite of PRI-724, blocks the binding of β -catenin to cAMP-

1
2
3
4
5
6 **response element-binding protein (CREB)-binding protein (CBP) and inhibits the**
7
8
9 **differentiation of lung fibroblasts to myofibroblasts induced by TGF- β 1.**

10
11
12
13 ICG-001, a compound similar to PRI-724, was reported to specifically disrupt the
14
15
16 interaction between β -catenin and CBP, not p300, in the nucleus (24). We therefore
17
18
19 analyzed the activity of C-82 to block the binding of β -catenin and CBP in lung fibroblast
20
21
22
23 B6 cells. B6 cells were stimulated with Wnt3a protein in the presence or absence of 1 or
24
25
26 2 μ M C-82. The cell lysate was harvested 12 hours after Wnt3a stimulation, and was
27
28
29 immunoprecipitated with anti-CBP or p300 antibody. The whole cell lysate and
30
31
32 immunoprecipitated samples were electrophoresed and plotted with anti-nonphospho- β -
33
34
35 catenin antibody. Treatment with C-82 clearly reduced the binding of β -catenin and CBP
36
37
38 (Figure 1C). On the contrary, the binding of p300 to β -catenin was increased after C-82
39
40
41 treatment. Next, we examined the biological effect of C-82 on the myofibroblast
42
43
44 differentiation of lung fibroblasts. Wnt/ β -catenin signaling was reported to play a critical
45
46
47 role in the TGF- β signaling pathway (25, 26). C-82 dose-dependently reduced the
48
49
50 expression of α -smooth muscle actin (SMA) in MRC5, a human lung fibroblast cell line,
51
52
53 treated with TGF- β 1 (Figure 1D, **Supplementary** Figure 4), indicating the inhibition of
54
55
56
57
58
59
60

1
2
3
4
5
6 myofibroblast differentiation.
7
8
9

10
11
12
13 **PRI-724 ameliorates bleomycin-induced lung fibrosis with a late, but not an early,**
14
15
16 **treatment schedule.**
17

18
19 We induced lung fibrosis in mice using the intratracheal administration of bleomycin.

20
21 Administration with PRI-724 alone did not induce any adverse events including weight
22
23 loss (data not shown) or fibrotic changes determined by the hydroxyproline content of the
24
25 lungs (Figure 2A), suggesting the absence of toxic effects of PRI-724. Next, antifibrotic
26
27 effects of PRI-724 were examined with an early treatment schedule (Figure 2B). In this
28
29 setting, bleomycin sulfate (1.25 mg/kg) was injected via the oropharyngeal aspiration on
30
31 day 0, and PRI-724 (1 or 10 mg/kg per day) or PBS was administered on day 3
32
33 subcutaneously using an osmotic pump. Three weeks after the bleomycin injection, we
34
35 sacrificed mice, and measured the collagen content. As shown in Figure 2B, there were
36
37 no significant differences in hydroxyproline levels among groups treated with bleomycin.
38
39

40
41 We next examined a late treatment schedule using PRI-724. Under this experimental
42
43 condition, the administration of PRI-724 was started on day 21, three weeks after the
44
45
46
47
48
49
50
51
52

1
2
3
4
5
6 bleomycin injection. After PRI-724 treatment for two weeks, the mice were sacrificed
7
8
9
10 five weeks after the bleomycin injection to examine the fibrotic changes in the lungs. The
11
12
13 hydroxyproline content in the PRI-724 group was significantly decreased compared with
14
15
16 that in the PBS group (393.4 ± 21.2 vs. 325.3 ± 17.3 $\mu\text{g}/\text{lung}$, $P < 0.05$) (Figure 2C). As
17
18
19 shown in Figure 3, the histological examination also showed that the fibrotic changes
20
21
22
23 around the peri-bronchiolar lesion in the lungs were reduced in the PRI-724 group.
24
25
26
27
28
29

30 **Immunofluorescence analysis of active β -catenin in various cells in the fibrotic lungs** 31 32 33 **of mice.** 34 35

36 We analyzed the expression of active β -catenin in various cells of the lungs treated with
37
38
39 bleomycin for three weeks in mice with double immunofluorescence staining. To detect
40
41
42 active β -catenin, antibodies for nonphospho- and pY654- β -catenin were used. As shown
43
44
45 in Figure 4A and 4B, alveolar macrophages detected by anti-CD68 antibody were
46
47
48 strongly stained with both anti-nonphospho- and pY654- β -catenin antibodies. In most
49
50
51 macrophages, their membranes were clearly stained, and the nucleus was also stained in
52
53
54 some cells (Figure 4A and 4B, Supplementary Figure 5A, 5B), indicating the activation
55
56
57
58
59
60

1
2
3
4
5
6 of β -catenin. Regarding AECs, nonphospho- β -catenin was detected by double-staining
7
8
9
10 with pro-SPC, but the level of staining was less than that seen in macrophages (Figure
11
12
13 4C, Supplementary Figure 5C). The nuclear localization of nonphospho- β -catenin was
14
15
16 also observed in some AECs (Figure 4C, Supplementary Figure 5C). The myofibroblasts
17
18
19 were stained with anti- α -SMA antibody. As shown in Figure 4D, positive staining with
20
21
22 anti-pY654- β -catenin antibody was detected in myofibroblasts, but the level was weak
23
24
25 compared with that in macrophages. It was difficult to determine the nuclear localization
26
27
28 of pY654- β -catenin in fibroblast-like cells. In addition, we found other cells, which were
29
30
31 positive for Sca-1, which strongly stained for pY654- β -catenin in the fibrotic lungs, and
32
33
34 some cells also showed nuclear localization (Figure 4E).
35
36
37
38
39
40
41
42

43 **Effects of PRI-724 in BAL cells in the fibrotic lungs of mice.**

44
45
46 To examine the effects of PRI-724 on the number and classification of BAL cells, mice
47
48
49 were sacrificed on day 28, one week after **starting** PRI-724 administration **with a late**
50
51
52 **treatment schedule**, and BAL was performed. As shown in Figure 5A, the total cell
53
54
55 number showed a decreasing tendency in the group receiving PRI-724 treatment,
56
57
58
59
60

1
2
3
4
5
6
7 although the difference was not significant. However, the number of macrophages was
8
9
10 significantly reduced in mice treated with PRI-724 compared with the PBS group ($8.3 \pm$
11
12
13 0.89 vs. $6.8 \pm 1.09 \times 10^4/\text{mL}$, respectively, $P < 0.05$) (Figure 5A).
14
15

16 17 18 19 20 **PRI-724 reduces the production of TGF- β 1 in bleomycin-induced lung fibrosis in** 21 22 23 **mice**

24
25
26 Next, we examined the level of active TGF- β 1 in BAL fluid one (day 28) and two (day
27
28
29 35) weeks after the administration of PRI-724, as assessed by ELISA. The level of active
30
31
32 TGF- β 1 in BAL fluid in the PRI-724 groups was decreased significantly on day 28, one
33
34 week after the PRI-724 treatment (Figure 5B). However, there was no significant
35
36
37 difference in the TGF- β 1 level on day 35, two weeks after the PRI-724 treatment,
38
39
40 although a decreasing tendency was observed (Figure 5B). We also analyzed the
41
42
43 expression level of TGF- β 1 in BAL fluid collected from mice treated with an early
44
45
46 treatment schedule as shown in Figure 2B. However, there was no significant difference
47
48
49 in the TGF- β 1 level on both day 7 and 14 (Supplementary Figure 6). These data support
50
51
52
53 that PRI-724 ameliorates bleomycin-induced lung fibrosis with **the** late, but not **the** early,
54
55
56
57
58
59
60

1
2
3
4
5
6
7 treatment schedule.
8
9

10 We finally examined the effects of C-82 on the production of TGF- β 1 by AMs. Murine
11
12
13 AMs were harvested from the BAL fluids one week after bleomycin injection. After 24
14
15
16 hours of treatment with C-82 or solvent in 1.0% FBS RPMI1640 medium, the level of
17
18
19 TGF- β 1 was measured in the supernatant treated with acid activation by ELISA, and the
20
21
22 mRNA level of cells by RT-PCR. As shown in Figure 6B, the level of latent TGF- β 1
23
24
25 protein was decreased in the C-82 group (Figure 6A), and the level of TGF- β 1 mRNA
26
27
28 was also suppressed in that group (Figure 6B).
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

DISCUSSION

In the present study, we examined the activated β -catenin signaling in various types of cells in the lungs treated with bleomycin by immunofluorescence staining, and found that AMs were stained most strongly, particularly in the nucleus, among lung cells including AECs and mesenchymal cells like fibroblasts. In addition, we demonstrated the antifibrotic effects of a novel inhibitor PRI-724, specifically disrupting the interaction of β -catenin and CBP, on pulmonary fibrosis induced by bleomycin in mice partly via inhibiting the production of TGF- β 1 by AMs.

The Wnt/ β -catenin signaling pathway was established to be highly activated in hyperplastic AECs and bronchiolar epithelial cells in IPF as well as in experimental pulmonary fibrosis model mice (7-11). However, as the specific deletion of β -catenin in AECs showed the deterioration of lung fibrosis and delayed recovery from lung injury, β -catenin signaling in AECs may serve rather to maintain the survival and promote regeneration in the fibrotic lungs (12), although profibrotic mediators such as WNT1-inducible signaling protein-1 was simultaneously induced (11). On the other hand, blockade of the Wnt/ β -catenin signaling pathway using inhibitors or inhibitory protein

1
2
3
4
5
6 Dickkopf-1 (DKK-1) suppressed pulmonary fibrosis in mice (14-18). These results
7
8
9 suggest that major target cells for Wnt/ β -catenin inhibition to inhibit pulmonary fibrosis
10
11
12 are cells other than AECs, such as fibroblasts or myeloid cells that have migrated into the
13
14
15 lungs. The fibroblasts could be possible targets because lung fibroblasts were reported to
16
17
18 proliferate via activated β -catenin signaling including canonical and TGF- β -mediated
19
20
21 pathways, and localized β -catenin staining of the nuclei of fibroblasts was observed (8,
22
23
24 9, 13, 26). In addition, Sennello et al recently reported that the activation of β -catenin
25
26
27 signaling in lung myeloid cells such as macrophages, neutrophils, B lymphocytes, and T
28
29
30 lymphocytes, except eosinophils, in the Axin2- β -galactosidase reporter mouse (19). They
31
32
33 also mentioned that the majority of β -catenin-activated cells appear morphologically to
34
35
36 be of immune cell origin, although there are some nonimmune cell types in the distal
37
38
39 lungs, such as AECs or fibroblasts (19). These observations are consistent with our results
40
41
42 of immunofluorescence staining of the fibrotic lungs, in which strong signals of activated
43
44
45 β -catenin were frequently observed in CD68-positive macrophages, but less in AECs or
46
47
48 fibroblasts. In addition, we also observed an activated β -catenin signal in Sca-1-positive
49
50
51 progenitor-like cells. Although these cells are CD45-negative and co-stained for TGF- β 1
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7 (data not shown), the role in pulmonary fibrosis should be explored in a further study.
8
9

10 Next, we investigated the antifibrotic effects of PRI-724, a novel second-generation-
11
12
13 specific inhibitor for the interaction between β -catenin and CBP (27). The first-generation
14
15
16 β -catenin/CBP inhibitor ICG-001 has been demonstrated to show antifibrotic effects on
17
18
19 various organs with fibrosis, including the lungs, skin, and liver (14, 28, 29). PRI-724 is
20
21
22 a novel inhibitor generated for clinical use (27). Notable points regarding the antifibrotic
23
24
25 effects of PRI-724 were the effective phase and mode of action in pulmonary fibrosis.
26
27
28

29 The antifibrotic effects were observed when treatment with PRI-724 was started on day
30
31
32 21 in the progressive and/or resolution phase, although the initiation of PRI-724 from day
33
34
35 3 in the inflammatory phase did not lead to any antifibrotic effects. These results may
36
37
38 indicate the therapeutic effects of PRI-724 because the effects in an early phase in a
39
40
41 bleomycin model were mediated predominantly by anti-inflammatory and not antifibrotic
42
43
44 effects. In addition, early treatment may induce dual bidirectional effects, profibrotic and
45
46
47 antifibrotic, which are mediated by AECs and fibroblasts in response to epithelial injury.
48
49
50

51 The mechanistic study identified a reduced number of AMs on day 28 one week after
52
53
54 PRI-724 administration. These results are consistent with the previous data in which the
55
56
57
58
59
60

1
2
3
4
5
6
7 number of monocyte-derived Siglec-F^{low} recruited AMs was decreased in bleomycin-
8
9
10 treated mice lacking lipoprotein receptor-related protein 5 (Lrp5), which is a Wnt
11
12
13 coreceptor and mediates a positive signal in the Wnt/ β -catenin pathway (19). These data
14
15
16 suggest that β -catenin signaling promotes the differentiation of recruited monocyte-AM
17
18
19 populations that contribute to the fibrotic phenotype (19). A more recent study reported
20
21
22 the involvement of the Wnt/ β -catenin signal in the proliferation of macrophages (30). In
23
24
25 addition to our immunofluorescence study, the Wnt/ β -catenin signal may be strongly
26
27
28 associated with activities of subpopulations of macrophages, particularly AMs. We also
29
30
31 identified a reduced level of TGF- β 1 in BALF on days 28, one week after PRI-724
32
33
34 initiation. This is also compatible with the results from Lrp^{-/-} mice, with a reduced level
35
36
37 of TGF- β production by alveolar leukocytes reported in the later, not early, phase of
38
39
40 pulmonary fibrosis (17). Furthermore, we noted the suppressive effects of C-82, an active
41
42
43 metabolite of PRI-724, on TGF- β 1 production of AMs *in vitro*. However, it was
44
45
46 demonstrated that the loss of Lrp5 and/or 6 does not have an impact on the TGF- β -
47
48
49 dependent up-regulation of TGF- β (17). There is a report that β -catenin is not involved
50
51
52 in Smad3-dependent transcriptional activities in the macrophage cell J774 (31). Thus, the
53
54
55
56
57
58
59
60

1
2
3
4
5
6 possibility that the reduction of TGF- β 1 expression is mediated by inhibition of the
7
8
9
10 Wnt/ β -catenin signal was supported by other reports (14-17), and another signaling
11
12
13 pathway interacting with CBP may be involved in the suppressive activity of PRI-724/C-
14
15
16 82 on AMs.

17
18
19 These results suggest that the novel β -catenin/CBP inhibitor PRI-724 is a potent
20
21
22 antifibrotic agent that acts through modulating the activity of AMs in the lungs, although
23
24
25 a further mechanistic study is required to explore the mode of action.
26
27
28
29
30
31
32

33 **Acknowledgements**

34
35
36 The authors thank Tomoko Oka and Megumi Kume for technical assistance. We also
37
38
39 thank the members of the Nishioka lab for their technical advice and fruitful discussions.
40
41
42
43
44
45

46 **Fundings**

47
48
49 This work was supported by the Japan Society for the Promotion of Science (JSPS)
50
51
52
53 KAKENHI under Grant (Number JP16H0530910), Japan Agency for Medical Research
54
55
56 and Development (AMED) under Grant, and the Ministry of Health, Labour and Welfare,
57
58
59
60

1
2
3
4
5
6
7 the Study Group on Diffuse Pulmonary Disorders, Scientific Research/Research on
8
9
10 Intractable Diseases under Grant (Number 0000025921).
11
12
13
14
15

16 **Declaration of interest statement**

17
18
19
20
21 Y.N. receives research funding from PRISM Pharma Co., Ltd.
22
23
24
25
26

27 **REFERENCES**

- 28
29
30
31 1. Richeldi L, Collard HR, Jones MG. Idiopathic pulmonary fibrosis. *Lancet*. 2017;
32
33 389: 1941-1952. DOI: 10.1016/S0140-6736(17)30866-8 [PubMed: 28365056]
34
35
36
37 2. Nishioka Y. Pathogenesis of IPF. Is abnormal repair of epithelial damage involved
38
39 in the basic pathogenesis of this disease? In: Nakamura H, Aoshiha K (eds)
40
41 Idiopathic Pulmonary Fibrosis, Springer, Tokyo. p43-58, 2016.
42
43
44
45
46
47 3. Raghu G, Rochweg B, Zhang Y, et al. American Thoracic Society; European
48
49 Respiratory society; Japanese Respiratory Society; Latin American Thoracic
50
51 Association. An Official ATS/ERS/JRS/ALAT Clinical Practice Guideline:
52
53
54
55
56
57
58 Treatment of Idiopathic Pulmonary Fibrosis. An Update of the 2011 Clinical
59
60

- 1
2
3
4
5
6
7 Practice Guideline. *Am J Respir Crit Care Med.* 2015; 192(2):e3-19. DOI:
8
9
10 10.1164/rccm.201506-1063ST [PubMed: 26177183]
11
12
13 4. Selman M, Pardo A, Kaminski N. Idiopathic pulmonary fibrosis: aberrant
14
15
16 recapitulation of developmental programs? *PLoS Med.* 2008; 5:e62. DOI:
17
18
19 10.1371/journal.pmed.0050062 [PubMed: 18318599]
20
21
22
23 5. Königshoff M, Eickelberg O. WNT signaling in lung disease: a failure or a
24
25
26 regeneration signal? *Am J Respir Cell Mol Biol.* 2010; 42:21-31. DOI:
27
28
29 10.1165/rcmb.2008-0485TR [PubMed: 19329555]
30
31
32
33 6. Guo Y, Xiao L, Sun L, Liu F. Wnt/ β -catenin signaling: a promising new target for
34
35
36 fibrosis diseases. *Physiol Res.* 2012; 61:337-46. [PubMed: 22670697]
37
38
39
40 7. Chilosi M, Poletti V, Zamò A, et al. Aberrant Wnt/beta-catenin pathway activation
41
42
43 in idiopathic pulmonary fibrosis. *Am J Pathol.* 2003; 162:1495-502. [PubMed:
44
45
46 12707032]
47
48
49
50 8. Xia H, Seeman J, Hong J, et al. Low $\alpha(2)\beta(1)$ integrin function enhances the
51
52
53 proliferation of fibroblasts from patients with idiopathic pulmonary fibrosis by
54
55
56 activation of the β -catenin pathway. *Am J Pathol.* 2012; 181:222-33. DOI:
57
58
59
60

1
2
3
4
5
6
7 10.1016/j.ajpath.2012.03.034 [PubMed: 22642910]
8
9

- 10 9. Königshoff M, Balsara N, Pfaff EM, et al. Functional WNT signaling is increased
11
12 in idiopathic pulmonary fibrosis. PLoS One. 2008; 3:e2142. DOI:
13
14

15
16 10.1371/journal.pone.0002142 [PubMed: 18478089]
17
18

- 19
20 10. Flozak AS, Lam AP, Russell S, et al. Beta-catenin/T-cell factor signaling is
21
22 activated during lung injury and promotes the survival and migration of alveolar
23
24 epithelial cells. J Biol Chem. 2010; 285:3157-67. DOI: 10.1074/jbc.M109.070326
25
26
27

28
29 [PubMed: 19933277]
30
31

- 32
33 11. Königshoff M, Kramer M, Balsara N, et al. WNT1-inducible signaling protein-1
34
35 mediates pulmonary fibrosis in mice and is upregulated in humans with idiopathic
36
37 pulmonary fibrosis. J Clin Invest. 2009; 119:772-87. DOI: 10.1172/JCI33950
38
39
40

41
42 [PubMed: 19287097]
43
44

- 45
46 12. Tanjore H, Degryse AL, Crossno PF, et al. β -catenin in the alveolar epithelium
47
48 protects from lung fibrosis after intratracheal bleomycin. Am J Respir Crit Care Med.
49
50

51
52 2013; 187:630-9. DOI: 10.1164/rccm.201205-0972OC [PubMed: 23306543]
53
54

- 55
56 13. Lam AP, Flozak AS, Russell S, et al. Nuclear β -catenin is increased in systemic
57
58

- 1
2
3
4
5
6
7 sclerosis pulmonary fibrosis and promotes lung fibroblast migration and
8
9
10 proliferation. *Am J Respir Cell Mol Biol.* 2011; 45:915-22. DOI:
11
12 10.1165/rcmb.2010-0113OC [PubMed: 21454805]
13
14
15
16 14. Henderson WR Jr, Chi EY, Ye X, et al. Inhibition of Wnt/beta-catenin/CREB
17
18 binding protein (CBP) signaling reverses pulmonary fibrosis. *Proc Natl Acad Sci*
19
20 USA. 2010; 107:14309-14. DOI: 10.1073/pnas.1001520107 [PubMed: 20660310]
21
22
23
24
25
26 15. Wang C, Zhu H, Sun Z, et al. Inhibition of Wnt/ β -catenin signaling promotes
27
28 epithelial differentiation of mesenchymal stem cells and repairs bleomycin-induced
29
30 lung injury. *Am J Physiol Cell Physiol.* 2014; 307(3): C234-44. DOI:
31
32 10.1152/ajpcell.00366.2013 [PubMed: 24898581]
33
34
35
36
37
38
39 16. Chen X, Shi C, Meng X, et al. Inhibition of Wnt/ β -catenin signaling suppresses
40
41
42 bleomycin-induced pulmonary fibrosis by attenuating the expression of TGF- β 1 and
43
44 FGF-2. *Exp Mol Pathol.* 2016; 101(1):22-30. DOI: 10.1016/j.yexmp.2016.04.003
45
46
47 [PubMed: 27112840]
48
49
50
51
52
53 17. Lam AP, Herazo-Maya JD, Sennello JA, et al. Wnt coreceptor Lrp5 is a driver of
54
55
56 idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med.* 2014; 190(2):185-95.
57
58
59
60

DOI: 10.1164/rccm.201401-0079OC [PubMed: 24921217]

18. Akhmetshina A, Palumbo K, Dees C, et al. Activation of canonical Wnt signalling is required for TGF- β -mediated fibrosis. *Nat Commun.* 2012; 3:735. DOI: 10.1038/ncomms1734 [PubMed: 22415826]
19. Sennello JA, Misharin AV, Flozak AS, et al. Lrp5/ β -catenin signaling controls lung macrophage differentiation and inhibits resolution of fibrosis. *Am J Respir Cell Mol Biol.* 2017; 56:191-201. DOI: 10.1165/rcmb.2016-0147OC [PubMed: 27668462]
20. Aono Y, Nishioka Y, Inayama M, et al. Imatinib as a novel antifibrotic agent in bleomycin-induced pulmonary fibrosis in mice. *Am J Respir Crit Care Med.* 2005;171(11):1279-85. DOI: 10.1164/rccm.200404-531OC [PubMed: 15735062]
21. Azuma M, Nishioka Y, Aono Y, et al. Role of alpha1-acid glycoprotein in therapeutic antifibrotic effects of imatinib with macrolides in mice. *Am J Respir Crit Care Med* 2007; 176:1243–1250. DOI: 10.1164/rccm.200702-178OC [PubMed:17717205]
22. Sato S, Shinohara S, Hayashi S, et al. Anti-fibrotic efficacy of nintedanib in pulmonary fibrosis via the inhibition of fibrocyte activity. *Respir Res.* 2017;

1
2
3
4
5
6
7 18(1):172. DOI: 10.1186/s12931-017-0654-2 [PubMed: 28915889]
8
9

10 23. Aono Y, Kishi M, Yokota Y, et al. Role of platelet-derived growth factor/platelet-
11
12
13 derived growth factor receptor axis in the trafficking of circulating fibrocytes in
14
15
16 pulmonary fibrosis. *Am J Respir Cell Mol Biol*. 2014; 51(6):793-801. DOI:
17
18
19 10.1165/rcmb.2013-0455OC [PubMed: 24885373]
20
21
22

23 24. Emami KH, Nguyen C, Ma H, et al. A small molecule inhibitor of beta-
24
25
26 catenin/CREB-binding protein transcription [corrected]. *Proc Natl Acad Sci U S A*.
27
28
29 2004; 101(34):12682-7 DOI: 10.1073/pnas.0404875101 [PubMed: 15314234]
30
31
32

33 25. Zhou B, Liu Y, Kahn M, et al. Interactions between β -catenin and transforming
34
35
36 growth factor- β signaling pathways mediate epithelial-mesenchymal transition and
37
38
39 are dependent on the transcriptional co-activator cAMP-response element-binding
40
41
42 protein (CREB)-binding protein (CBP). *J Biol Chem*. 2012; 287(10):7026-38. DOI:
43
44
45 10.1074/jbc.M111.276311 [PubMed: 22241478]
46
47
48

49 26. Xu L, Cui WH, Zhou WC, et al. Activation of Wnt/ β -catenin signalling is required
50
51
52 for TGF- β /Smad2/3 signalling during myofibroblast proliferation. *J Cell Mol Med*.
53
54
55 2017; 21(8):1545-1554. DOI: 10.1111/jcmm.13085 [PubMed: 28244647]
56
57
58
59
60

- 1
2
3
4
5
6
7 27. Tokunaga Y, Osawa Y, Ohtsuki T, et al. Selective inhibitor of Wnt/ β -catenin/CBP
8
9
10 signaling ameliorates hepatitis C virus-induced liver fibrosis in mouse model. *Sci*
11
12
13 *Rep.* 2017; 7(1):325. DOI: 10.1038/s41598-017-00282-w [PubMed: 28336942]
14
15
- 16 28. Beyer C, Reichert H, Akan H, et al. Blockade of canonical Wnt signalling
17
18 ameliorates experimental dermal fibrosis. *Ann Rheum Dis.* 2013; 72(7):1255-8.
19
20
21 DOI:10.1136/annrheumdis-2012-202544 [PubMed: 23595143]
22
23
24
- 25 29. Akcora BÖ, Storm G, Bansal R. Inhibition of canonical WNT signaling pathway
26
27 by β -catenin/CBP inhibitor ICG-001 ameliorates liver fibrosis in vivo through
28
29
30 suppression of stromal CXCL12. *Biochim Biophys Acta.* 2018; 1864(3):804-818.
31
32
33
34
35
36
37 DOI: 10.1016/j.bbadis.2017.12.001 [PubMed: 29217140]
38
39
- 40 30. Feng Y, Liang Y, Ren J, Dai C. Canonical Wnt Signaling Promotes Macrophage
41
42
43 Proliferation during Kidney Fibrosis. *Kidney Dis (Basel).* 2018; 4(2):95-103. DOI:
44
45
46
47 10.1159/000488984 [PubMed: 29998124]
48
49
- 50 31. Tian X, Zhang J, Tan TK, et al. Association of β -catenin with P-Smad3 but not
51
52
53 LEF-1 dissociates in vitro profibrotic from anti-inflammatory effects of TGF- β 1. *J*
54
55
56
57 *Cell Sci.* 2013; 126(Pt 1):67-76. DOI: 10.1242/jcs.103036 [PubMed: 23203799]
58
59
60

FIGURE LEGENDS

Figure 1. Activation of β -catenin in lung fibroblasts and macrophages and effects of C-82 on the binding of β -catenin with CBP and TGF- β -induced differentiation of lung fibroblasts. A: Lung fibroblast B6 cells were stimulated with Wnt3a (50 ng/mL) for 1 hour. The accumulation of β -catenin in the nuclei was examined with anti-nonphospho (active) β -catenin antibody. B: The accumulation of β -catenin in the nuclei in the macrophage cell line RAW257.6 was examined after treatment with Wnt3a for 12 hours with anti-nonphospho (active)- β -catenin antibody. C: The effects of C-82 on the interaction between β -catenin and CBP were examined with the nuclear extracts of lung fibroblast B6 cells stimulated with Wnt3a for 90 min. D: Effects of C-82 on the expression of α -SMA in MRC5, a human lung fibroblast cell line. MRC5 cells were cultured with TGF- β 1 (5 ng/mL) in the presence or absence of various concentrations of C-82 for 48 hours. The expression of α -SMA was examined with Western blotting. Data are representative of two separate experiments. The relative expression of each band was calculated by densitometric analysis. Data were analyzed using a Student's t-test and

1
2
3
4
5
6 displayed as means \pm SE. * $P < 0.05$ versus the control group in Figure 1A and 1B, and
7
8
9
10 versus the TGF- β only group in Figure 1D.

11
12
13
14
15
16 **Figure 2. PRI-724 reduces pulmonary fibrosis with a late treatment schedule in a**
17
18
19
20 **bleomycin-induced pulmonary fibrosis model.** A: C57BL/6 mice were treated with the
21
22 subcutaneous injection of PBS or PRI-724 (1 or 10 mg/kg per day) with osmotic
23
24 minipumps from days 3 to 21, and left lobe was harvested for analysis of the collagen
25
26
27 content by the hydroxyproline assay on day 21. B: C57BL/6 mice were treated with the
28
29 intratracheal administration of saline or bleomycin on day 0. In bleomycin groups, mice
30
31 were injected subcutaneously with osmotic minipumps containing PBS or PRI-724 (1 or
32
33 10 mg/kg/day) from days 3 to 21, and the left lobe was harvested for analysis of the
34
35 collagen content by hydroxyproline assay on day 21. C: C57BL/6 mice were treated with
36
37 the intratracheal administration of saline or bleomycin on day 0. In bleomycin groups,
38
39 mice were injected subcutaneously with osmotic minipumps containing PBS or PRI-724
40
41 (10 mg/kg/day) from days 21 to 35, and whole lungs were harvested for analysis of the
42
43 collagen content by the hydroxyproline assay on day 35. Data are presented as the mean
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7 \pm SE. (A) n=4; (B) n=6-9; (C) n=7-9, $P < 0.05$ (Student's t-test)
8
9
10
11
12

13 **Figure 3. Histological evaluation of the anti-fibrotic effects of PRI-724 on mice with**
14 **bleomycin-induced lung fibrosis.** C57BL/6 mice were treated with the intratracheal
15
16 bleomycin-induced lung fibrosis. C57BL/6 mice were treated with the intratracheal
17 administration of saline or bleomycin on day 0. In bleomycin groups, mice were injected
18
19 subcutaneously with osmotic minipumps containing PBS or PRI-724 (10 mg/kg/day)
20
21 from days 21 to 35, and the right lobe was harvested for histological examination on day
22
23 35. Hematoxylin and eosin (H&E) (A, C, and E) and azan mallory (B, D, and F) (scale
24
25 bar = 100 μ m) stainings were performed. (A, B) Saline. (C, D) BLM and PBS. (E, F)
26
27 BLM and PRI-724 (10 mg/kg).
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 **Figure 4. Immunofluorescence images of the β -catenin-positive cells' nuclei,**
44 **membranes, and cytoplasm in murine lungs treated with bleomycin.**
45
46
47

48
49 Immunofluorescent analyses were performed in the murine lungs on day 21 after the
50
51 bleomycin (1.5 mg/kg) injection. β -Catenin was stained with anti-phospho-Y654- β -
52
53 catenin (A, D, E) or nonphospho (active)- β -catenin (B, C) antibodies. Macrophages were
54
55
56
57
58
59
60

1
2
3
4
5
6 also stained with anti-CD68 antibody (A, B). Alveolar epithelial cells were stained with
7
8
9 anti-pro-SPC-antibody (C). Lung myofibroblasts were stained with anti- α -smooth
10
11
12 muscle actin antibody (D). Sca-1-positive cells were analyzed (E). Arrows indicate the
13
14
15
16 nuclear staining of each β -catenin. Scale bars, 10 μ m.
17
18
19
20
21
22


23 **Figure 5. Analysis of bronchoalveolar lavage fluid.** BAL was performed on day 28 or
24
25
26 35 after one or two weeks of treatment with PRI-724. A: Cell classification of BAL cells
27
28
29 on day 28 (n=6: BLM/PBS, n=5: BLM/PRI-724). B: Level of TGF- β 1 in BAL fluids on
30
31
32 days 28 and 35. Active TGF- β 1 was measured with ELISA (n=6: saline/PBS/day 28,
33
34
35 n=14: BLM/PBS/day28, n=12: BLM/PRI-724/day 28, n=6: BLM/PBS/day 35, n=6:
36
37
38 BLM/PRI-724/day 35). Data are the mean \pm SE. $P < 0.05$ using Student's t-test.
39
40
41
42
43
44
45
46

47 **Figure 6. C-82 inhibits the expression of TGF- β 1 on alveolar macrophages.** The AMs
48
49
50 were collected from mice 1 week after treatment with bleomycin with the BAL procedure.
51
52
53 The AMs were treated with saline or C-82 (1 μ M) in 1.0% FBS RPMI1640 medium for
54
55
56 24 hours, and the supernatants were collected. The **latent** TGF- β 1 levels were measured
57
58
59
60

1
2
3
4
5
6 using ELISA (n=4 in each group) (A). Real-time PCR was performed using RNA
7
8
9
10 extracted from the AMs after being treated with DMSO or C-82 for 24 hours (n=4 in each
11
12
13 group) (B). Data are the mean \pm SE. $P < 0.05$ using Student's t-test.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Fig. 1










Fig.3



H&E staining

Azan staining



saline



BLM/
PBS



BLM/
PRI-724
(10 mg/kg)







Fig.5

A



B






Fig.6

A



B



1
2
3
4
5
6
7 The novel inhibitor PRI-724 for Wnt/ β -catenin/CBP signaling ameliorates
8
9
10 bleomycin-induced pulmonary fibrosis in mice
11
12
13
14
15

16 Hiroyasu Okazaki¹, Seidai Sato¹, Kazuya Koyama¹, Shun Morizumi¹, Shuichi Abe¹,
17
18 Momoyo Azuma¹, Yajuan Chen¹, Hisatsugu Goto¹, Yoshinori Aono^{1,2}, Hirohisa Ogawa³,
19
20
21
22 Kozo Kagawa¹, Haruka Nishimura¹, Hiroshi Kawano¹, Yuko Toyoda¹, Hisanori Uehara³,
23
24
25
26
27 Hiroyuki Kouji^{4,5}, Yasuhiko Nishioka¹
28
29
30
31

32 **SUPPLEMENT DATA**
33
34
35

36 **FIGURE LEGENDS**
37
38
39

40 **Supplementary Figure 1. Expanded blots from Figure 1A**
41
42

43 **Supplementary Figure 2. Expanded blots from Figure 1B**
44
45

46 **Supplementary Figure 3. Activation of β -catenin in macrophages and effects of C-**
47
48
49
50 **82 on the binding of β -catenin with CBP.**
51

52
53 The accumulation of β -catenin in the nuclei in the macrophage cell line RAW257.6 was
54
55
56 examined after treatment with Wnt3a for 12 hours with pY654- β -catenin antibodies.
57
58
59
60

1
2
3
4
5
6
7 **Supplementary Figure 4. Expanded blots from Figure 1D**
8
9

10 **Supplementary Figure 5. Low magnification images of Figure 4.**
11
12




13 **Immunofluorescence** analyses were performed in the murine lungs on day 21 after the
14
15
16 bleomycin (1.5 mg/kg) injection. β -Catenin was stained with anti-pY654- β -catenin (A)
17
18 or nonphospho (active)- β -catenin (B, C) antibodies. Macrophages were stained with anti-
19
20 CD68 antibody (A, B). Alveolar epithelial cells were stained with anti-pro-SPC-antibody
21
22 (C). Arrows indicate the nuclear staining of each β -catenin. Scale bars, 100 μ m.
23
24
25
26
27
28
29

30 **Supplementary Figure 6. The expression level of TGF- β in BAL fluid collected from**
31
32
33 **mice treated with an early treatment schedule.**
34
35

36 C57BL/6 mice were treated with the intratracheal administration of saline or bleomycin
37
38 on day 0. In bleomycin groups, mice were injected subcutaneously with osmotic
39
40 minipumps containing PBS or PRI-724 (1 or 10 mg/kg/day) started from days 3. BAL
41
42 was performed on day 7 or 14 with mice treated with an early treatment schedule. **Total**
43
44 TGF- β 1 was measured with ELISA (n=5 in each group). Data are the mean \pm SE. $P <$
45
46
47
48
49
50
51
52
53 0.05 using Student's t-test.
54
55
56
57
58
59
60




Supplementary Figure 1.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Supplementary Figure 2.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60





Supplementary Figure 3.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Supplementary Figure 5.


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

A




CD68 (red)
Phospho-Y654-β-catenin (green)

B




CD68 (red)
Nonphospho-β-catenin (green)

C



Pro-SPC (red)
Nonphospho-β-catenin (green)

Supplementary Figure 6.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60